

Breastfeeding, bonding, and olfaction: unlocking the potential of mother-infant odour exchange



S. Craig Roberts,^{a,b,*} Fabrice Damon,^c Karine Durand,^c Jan Havlíček,^d Dimitrios Kourtis,^a Ben Langford,^e Agnieszka Sorokowska,^f Piotr Sorokowski,^f Vivien Swanson,^a Jonathan Williams,^g Tatjana Arnoldi-Meadows,^g Dylan Brimaud,^c Daniela Dlouhá,^h Lucie Jelínková,^d Šárka Kaňková,^h Lenka Kapicová,^d Alice C. Poirier,^c Kateřina Roberts,^a Justus C. Sander,^g Dagmar Schwambergová,^d Michaela Silvestri,^a Nijing Wang,^g Emmanuel Simon,ⁱ and Benoist Schaal^f

^aSchool of Psychology, University of Stirling, Stirling, UK

^bBeing Human Lab, University of Wrocław, Wrocław, Poland

^cDevelopment of Olfactory Communication and Cognition Laboratory, Université Bourgogne Europe, Institut Agro, CNRS, INRAE, UMR CSGA, 21000 Dijon, France

^dDepartment of Zoology, Faculty of Science, Charles University, Prague, Czech Republic

^eUK Centre for Ecology & Hydrology, Penicuik, UK

^fInstitute of Psychology, University of Wrocław, Wrocław, Poland

^gMax Planck Institute for Chemistry, Mainz, Germany

^hDepartment of Philosophy and History of Science, Faculty of Science, Charles University, Prague, Czech Republic

ⁱObstetrics and Foetal Medicine, Université Bourgogne Europe, Dijon, France

Summary

Breastfeeding is crucial for infant survival, growth, and health, and it enhances maternal-infant bonding and well-being. However, breastfeeding rates typically fall below international targets, partly due to a high prevalence of latching difficulties, intermittent sucking, refusing the breast, or poor milk supply. Here, we propose that such uniquely human difficulties might be ameliorated by recognising, understanding, and facilitating the olfactory mechanisms that, in other mammals, regulate breastfeeding initiation and maternal–infant relationships in the first weeks of life. We briefly review evidence that odour mediates nipple-searching and suckling behaviour in other species, summarise the comparable evidence in humans, and outline pathways that could potentially reap hitherto unrealised benefits of olfactory communication between human mothers and neonates. We argue that enhanced awareness of such odour exchange could inform and enable changes in both policy and practice that might improve breastfeeding success and maternal-infant bonding, ultimately contributing to reduced infant mortality worldwide.

eBioMedicine
2026;123: 106086

Published Online 20
December 2025
<https://doi.org/10.1016/j.ebiom.2025.106086>

Copyright © 2025 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Keywords: Olfactory communication; Smell; Attachment; Skin-to-skin; Breastfeeding; Perinatal care

Introduction

The benefits of breastfeeding are widespread and well-known. For infants, breastfeeding supports survival and long-term health outcomes, influencing immunity, metabolism, neurocognitive development, and mental well-being. For mothers, it plays a crucial role in bonding and recovery after childbirth while also reducing the risk of breast cancer. Recent estimates suggest that increasing breastfeeding rates could prevent 823,000 annual deaths in children under five and 20,000 annual deaths from breast cancer.¹ Nonetheless, fewer than half of babies worldwide are breastfed within the first hour after birth or exclusively breastfed for the recommended six months,² both key World Health Organization (WHO) and UNICEF targets.

Difficulties with breastfeeding initiation (e.g. latching, poor sucking, or refusing the breast) are pervasive even when mothers intend to breastfeed: a third³ to half⁴ have difficulties on the first day. Such difficulties may cause fatigue, pain, loss of confidence, and distress in mothers.⁵ For infants, they can pose clinical risks, lead to excessive weight loss,^{3,4} and deprive them of the earliest opportunity to ingest colostrum and non-pathogenic microbiota from the breast. Such consequences may be especially perilous in medically underdeveloped regions and may have been responsible for countless deaths in previous centuries.⁶

However, finding and latching successfully onto the breast is by no means a uniquely human challenge. By definition, it confronts neonates of every mammalian species. A universal solution—found in all female mammals studied to date—is the production of chemosensory signals from the mammary glands (or immediately adjacent sources). These odour messages

*Corresponding author. School of Psychology, University of Stirling, Stirling, FK9 4LA, UK.

E-mail address: craig.roberts@stir.ac.uk (S.C. Roberts).

take advantage of neonatal capabilities in olfaction, which often mature earlier than audition and vision.^{7,8}

In this personal view paper, we propose that some of the difficulties with human breastfeeding might be ameliorated by recognising, understanding, and facilitating olfactory processes that underpin nipple-finding, latching, and sucking, especially in the critical first days of life. As in other social interactions in humans, these olfactory mechanisms have been widely underestimated and underappreciated. We advocate for a comprehensive reassessment of the importance of olfaction in mother–newborn interactions and describe how existing and developing knowledge of such processes could potentially be applied to inform systematic and transformational change in clinical practices around the mother and neonate that could have a significant positive impact on the initiation and subsequent maintenance of breastfeeding.

The need for such a reassessment is perhaps best illustrated by the fact that olfaction is not mentioned in the headline documents underpinning the WHO/UNICEF's Baby-Friendly Hospital Initiative (BFHI, revised in 2018), which aims to encourage the support of breastfeeding worldwide. Specifically, there is no explicit reference to odour or olfaction in its *Ten Steps to Successful Breastfeeding*, which outlines critical management procedures and clinical practices to promote optimal care for new mothers and infants, nor in the 56 pages of guidance provided for their implementation.⁹ However, as we will describe, a focused and explicit consideration of olfaction's role in the early stages of breastfeeding could have direct implications for each of the *Ten Steps*.

The myth of poor human olfaction

The surprising omission of references to odour and olfaction in key policy and implementation documents may stem from the longstanding perception that olfaction is unimportant in humans. Research has now undermined this view to the point of it being termed 'a 19th-century myth'.¹⁰ In fact, humans retain a neural architecture enabling us to outperform even 'super-smeller species' (dogs, mice) in detecting certain odours,¹⁰ and olfaction is increasingly recognised as playing a key role in our interactions with our environment.¹¹ We use it in food selection and flavour perception, detecting environmental hazards and pathogens, and social communication.

Regarding the latter, natural (or even perfumed) body odours convey characteristics such as identity, age, gender, fertility, and affective state, mediating processes such as attraction, familiarity, and attachment.¹² Yet, it has been repeatedly argued^{7,8,13,14} that these functions are overshadowed by the importance of olfaction in the critical maternal–infant relationship formed in the first days of life. As newborn infants are at their most vulnerable and otherwise sensorily limited, this is a

unique moment when odours can communicate life-saving information.

Across species, nipples convey a conspicuous 'smellscape'

Indeed, every mammalian neonate urgently needs to find and latch onto a nipple or teat to receive nourishment and the full benefit of the protective properties of colostrum. Over evolutionary time, therefore, natural selection has apparently favoured the conspicuousness of the nipple.⁷ This conspicuousness need not be visual; indeed, the production of olfactory 'beacons' from the mammary glands appears to be a universal feature of female mammals.⁷

How do we know that the sense of smell is involved? In various species, behavioural evidence includes positive orientation of neonates towards swabs rubbed over the mammary area compared to swabs with saline solution¹⁵ and towards the odour of lactating over virgin females.¹⁶ Nipple search behaviour is hindered or eliminated by nostril obstruction or rendering neonates anosmic,¹⁷ and latencies to find the nipple and begin sucking are increased by washing nipples.¹⁸

However, we know most about odour-guided neonatal behaviour in the European rabbit, *Oryctolagus cuniculus*, in which selection pressure on nipple-finding is intense because pups are born 'blind' and mothers visit the nesting burrow only once daily, and then only for 3–5 min. This pressure apparently led to the evolution of the compound 2-methyl-2-butenal in rabbit milk, now known as the rabbit mammary pheromone,¹⁹ having met the criteria required to satisfy the formal definition of a pheromone.¹³ Such criteria include that it releases unambiguous and functional behaviour (in this case, searching, grasping, and sucking; Fig. 1), that the same behaviour is not elicited by other reference compounds or in other species, and that the responses should not (or only minimally) depend on previous exposure and learning. The pheromone is present at its highest concentrations immediately postpartum, declines during lactation, and is undetectable by weaning; this pattern is matched by pups' responsiveness, as neonates react more strongly to it than pre-weanlings.²⁰ Pups that are naturally non-reactive to the mammary pheromone on day 1 die before weaning, demonstrating the consequences of natural selection in action.²¹ As in other species, latching is delayed by washing the abdominal fur of rabbit does or covering nipples with airtight rubber or film,^{22,23} and rendering pups anosmic leaves them unable to find the nipple.²³

Odour influences nipple-searching behaviour in infant humans

Although human mothers may bring their babies to the breast, human and rabbit infants share similar sensory




Fig. 1: Responses to odours soon after birth. On the left, a rabbit pup (a) searches and (b) grasps a glass rod when it is coated in the rabbit mammary pheromone. On the right, a 3-day-old infant shows (c) lip pursing and (d) tongue protrusion in response to Montgomery's gland secretion (Photo credits: CSGA-CNRS).

abilities and vulnerabilities, and there is strong evidence that olfaction similarly plays a key role in breastfeeding initiation and success in humans.⁸ Infant rooting (bilateral head movements and oral searching for the nipple²⁴) and mouthing behaviour²⁵ is stimulated by elements of the maternal smellscape.²⁶ Within minutes after birth, newborns positioned prone on the mother's abdomen can root and crawl independently towards the mother's breast,^{27,28} and breast odour is sufficient to elicit or enhance this movement.^{29–31} It also comforts infants, reducing crying²⁶ and expressions of stress and pain.³² If the lactating breast is washed, they are less likely and take longer to find it.²⁹ When the breast is covered with a scentless plastic film, infants are less attracted to it, exhibit fewer mouthing responses (Fig. 1), and are quicker to cry.²⁶

It cannot be ruled out that learning of breast odour occurs in utero or immediately after birth, since babies can learn odours in amniotic fluid³³ and within a few

hours postpartum,³⁴ and there are similarities in the chemical profiles of amniotic fluid and breast odour.³⁵ Nonetheless, postnatal learning is not necessarily involved in the very earliest responses.³⁶ Even infants who have been exclusively fed formula from birth orient their heads towards the breast odour of an unfamiliar, lactating woman more than the odour of their familiar formula^{24,37} and the breast odour of a non-lactating woman³⁸; they also suck harder and more frequently when a mother's milk odour is presented under their noses when feeding compared with the odour of formula or water.³⁹

Sources of breast odour active components

As in rabbits, milk is a likely source of the active chemical compounds responsible for newborn attraction to human breast odour. At least 63 volatile compounds in breastmilk are detectable by the human

nose.⁴⁰ However, different volatiles are expressed in lacteal secretion at different times (this is likely a general feature of all mammals⁴¹). When compounds were compared across three lactational stages (colostrum, transitional milk, mature milk), some occurred at relatively higher abundance in colostrum⁴² and might, therefore, be critical in facilitating nipple-finding and breastfeeding initiation. Indeed, behavioural experiments indicate that colostrum is especially attractive to 2-day-old newborns compared to mature milk.⁴³

Another potential source is skin glands surrounding the nipple, especially the Montgomery's glands (MGs, also known as areolar glands). Histological evidence shows these small protuberances combine both miniature mammary acini and sebaceous glands, the latter being involved in scent communication across taxa.⁷ Secretory MGs are unevenly distributed on the areola, being more common in the upper lateral quarter on each breast; such a distribution is indicative of communicatory function, as these quarters are those to which the feeding infant's nose is usually directed.⁴⁴ The number of MGs on the breast of individual women varies, with greater numbers predicting faster re-gain of initial infant body mass, earlier onset of lactation, faster latching onto the nipple, and higher sucking intensity.⁴⁴ Furthermore, the odour of MG secretion from an unfamiliar mother stimulates stronger oro-cephalic responses (lip-pursing, tongue protrusion; Fig. 1) in 3-day-old infants than an unfamiliar mother's milk, cow milk, or formula.⁴⁵

Odours influence maternal-infant bonding

So far, we have described maternal chemical signals (or chemosignals) that have evolved to communicate vital information to newborns; however, infant-produced odours may also alter maternal behaviour. In sheep, for example, ewes deprived of olfactory (but not auditory) cues from lambs show reduced care behaviour.⁴⁶ In rats, production of the compound dodecyl propionate by the pup's preputial gland stimulates anogenital licking by the mother, which is critical for pup survival through stimulating excretory reflexes.⁴⁷

In humans, infant odours appear to play a significant role in maternal bonding. The smell of a newborn baby is salient and attractive to post-parturient women⁴⁸ and activates reward-related neural substrates,⁴⁹ especially the odour of their own child.⁵⁰ Mothers rapidly learn to distinguish their baby's smell³¹; this ability, and how much they express liking their infant's odour, predicts subjective ratings of maternal bonding⁴⁸ and investment.⁵¹

Beyond their role in bonding, one can also speculate that the same evolutionary selection pressures that shape breast signals targeted at infants could lead to infant-produced chemosignals that communicate hunger or emotional state to caregivers. In rats, the odour of

hungry individuals is known to elicit caring behaviours.⁵² In adult humans, hunger or satiety can be discriminated by smell,⁵³ as can emotional states (e.g. fear, stress).⁵⁴ Such findings raise the possibility that infant odours could communicate hunger or distress, acting as a signal of need, analogous to an 'olfactory cry', perhaps in advance of an auditory one. Even small uplifts in maternal attentiveness arising from such a signal could have life-saving effects, perhaps especially at night when vision is ineffective. Such uplifts might occur subconsciously, as sub-threshold odours can affect behaviour in other contexts.⁵⁵ We can even hypothesise that increasing expression of such a chemosignal as infants become hungry could trigger the mother's physiological pathways that lead to milk letdown and/or upregulation in the emission of the volatile compounds comprising the breast odour signal.

Harnessing the benefits of odour communication in perinatal care

There are several possible pathways by which emerging insights into maternal-infant olfactory communication in humans and other mammals could be harnessed to boost the success of practical initiatives, such as those aimed at increasing breastfeeding rates (Fig. 2). More research is needed, but the existing evidence is already sufficient to suggest that we should aim to boost practices that favour olfactory communication to establish and maintain breastfeeding and discourage those that hinder it. Further translational opportunities await the discoveries of the precise chemical mechanisms involved.

Facilitating communicative mechanisms

The most obvious approach to boosting olfactory communication is maintaining mother-infant proximity in the earliest moments of postnatal interaction. Thus, providing rooming-in and skin-to-skin contact (hereafter, SSC)—both incorporated into the WHO/UNICEF's *Ten Steps*—are essential to enable communication in hospital settings (rooming-in does not apply when mothers give birth at home, as is increasingly common in some countries). Although the prevalence of SSC within the first hour of life varies hugely across the world, research demonstrates its well-known and varied benefits.⁵⁶ In infants, for example, it is associated with reduced stress and crying, and increased rates of breastfeeding initiation and exclusive breastfeeding. For mothers, SSC is associated with reduced stress and increased breastfeeding self-efficacy, as well as the release of oxytocin, which enhances parental bonding and care behaviours, and boosts lactation itself.

Even in clinical settings that value SSC, the sensory mechanisms underpinning SSC benefits are often either not explicitly considered or assumed to be




Fig. 2: Opportunities for integrating understanding and awareness of maternal-infant olfactory communication in clinical practice. *Above:* Odour management measures in perinatal care can address barriers to odour communication or act to facilitate them. *Below:* Identifying the specific chemical compounds responsible for eliciting adaptive behaviours, including the chemosignal that attracts newborns to the nipple and a (hypothesised) signal of need/distress from infants, could lead to the development of isolated or synthetic chemosignals with defined practical applications.

tactile.⁵⁷ However, olfaction is likely to be equally or even more involved.^{8,58,59} At least four of Widström’s nine critical behavioural phases²⁸ during SSC in the first hour involve smell (stages 4 and 6–8: activity, crawling towards the breasts, familiarisation with the areola/nipple, and initiation of suckling).^{28,60} Appreciating that the main benefit of SSC might come via olfaction—at least as much as touch or vision, and perhaps interacting with them multimodally—is essential to shaping clinical practices that facilitate it.

Furthermore, there is evidence to suggest that providing newborns with the opportunity to use their sense of smell in rooting and crawling to the breast is an important factor influencing breastfeeding outcomes. Klaus⁶¹ noted that an infant who is permitted to crawl unaided to the breast “opens his mouth widely and, after several attempts, makes a perfect placement on the areola of the nipple”, while in those placed directly between the breasts “the infant reaches the

target just as effectively, but with a different pattern of behaviour. Instead of beginning to suckle, many infants just lick the nipple and their hands” (p.1244). Righard and Alade²⁷ reported that infants allowed to crawl unaided were more likely to begin breastfeeding and less likely to suck inefficiently or with a superficial latch, compared with those placed directly at the breast and whose mothers usually manipulated the nipple into the infant’s mouth. A randomised controlled trial (RCT) recently showed that the crawl (facilitated by cheek-to-cheek contact and smearing the breast with expressed milk to enhance odour cues) led to reduced time to breastfeeding initiation and improved latching compared to SSC alone.⁶² While more studies are desirable, such findings suggest that active rooting and crawling provide opportunities for learning the mother’s breast and body odour and how to respond appropriately. It might even form a critical sensitive stage⁵⁹ (similar to that suggested for maternal

bonding^{63,64}), since subsequent recognition of maternal odour is improved in infants provided with early SSC⁶⁵ and neural activation in response to the odour of colostrum is higher in the first 24 h than over subsequent days.⁶⁶ Thus, the direct placement of a passive newborn on the breast may be (at first sight, counter-intuitively) less beneficial, and have long-lasting effects on infants, compared to providing them with a reasonable and appropriately timed opportunity to actively and independently find the breast, even if this means they take longer to initiate feeding.

Beyond SSC, further research is also needed on interventions to enhance communication, particularly where direct communication is not possible. To date, most work has focused on premature births, where mothers and newborns are often physically separated. Welch et al.⁶⁷ developed a 'family nurture intervention' (FNI) to counteract the adverse effects of separation in preterm families; among other intervention sub-components, this included a protocol involving reciprocal exchange between mothers and infants of cotton cloths infused with body odour (worn in the bra or placed under/on the newborn's head). The FNI had several positive outcomes in terms of mother-infant engagement (touch, gaze),⁶⁸ neurodevelopmental functioning (including cognition, language, attention, and social functioning up to 18 months, and theory-of-mind ability at age 4), and reduced rates of post-discharge anxiety and depression.⁶⁹ Odour exchange is an intuitive approach, and parents readily grasp its potential benefits: sometimes even parents in the standard care group wanted to use it,⁷⁰ and parents in one trial were more likely to engage with it than SSC, reporting that it enhanced feelings of attachment with the baby.⁷¹ However, it should be noted that the FNI is multi-modal, and more work is needed to isolate effects of scent exchange from other subcomponents (e.g. soothing voice, touch, SSC).

More separable approaches include the use of maternal odour to facilitate feeding or relieve pain. One meta-analysis of six RCTs⁷² indicated that odour stimulation with breastmilk reduces the transition time from tube to oral feeding; in these studies, the preterm infant's own mother's breastmilk was applied to gauze pads held 2–5 cm from the infant's nose while tube-feeding. However, another meta-analysis of eight studies found no significant benefit,⁷³ so further research is required on this issue. One study suggested a reduced likelihood of neurosensory impairment in premature infants exposed to the smell and taste of milk before gastric tube feeds.⁷⁴ Finally, a meta-analysis of eight RCTs measuring pain during clinical interventions (e.g. venipuncture, heel-prick) concluded that, compared to no-odour controls, infants exposed to maternal odour before or during the intervention have

lower pain scores and cry less.⁷⁵ Odour presentation likely reduces pain either by influencing affect (such as diminishing fearfulness⁷⁶) or by distraction.⁷⁷ Overall, these studies demonstrate that odour exchange interventions could be effective in achieving multiple desirable outcomes for mothers and infants in clinical care.

Removing barriers to odour communication

Clinical care practices that reduce the potential for effective odour transmission between mothers and infants should, as far as possible, be avoided. Perhaps the most obvious example is bathing newborns. Although the WHO recommends delaying the first bath for at least 24 h, bathing within 10 min of birth remains common in many countries.⁷⁸ Not only will this interfere with SSC and autonomous breastfeeding initiation by the infant if bathing happens before the first breastfeed,⁶¹ it also disrupts maternal reception of infant odours during the same period. Similarly (but less often discussed), reducing the frequency of breast-washing in the first days may ensure that critical olfactory cues are maximally available for reception by infants.²⁹

Physical barriers have been reduced since the broad adoption of the practice of rooming-in. However, the WHO's recommendation that babies have more layers of clothing than adults, and to keep the head covered with a hat, maintains a remaining physical barrier for the reception of infant odour by mothers; adjusting this guidance might be appropriate where thermoregulatory conditions and cultural traditions permit.

Furthermore, the introduction of olfactory interference in the birthing and early postnatal environment should be avoided. Competing olfactory practices might include the use of odour-rich antiseptic solutions,⁷⁹ the hand odours of practitioners touching the mother's breasts,⁶⁰ and perfume use by those around the mother. Odour-neutral disinfectants should likely be used where birthing and first breastfeeding episodes will occur, and it should be recognised that aromatherapy practices for pain relief in the second or third stages of labour⁸⁰ could produce unwelcome olfactory interference.

Recommendations concerning mothers' skincare or cosmetic use may be more nuanced. Ointments for sore or cracked nipples should ideally be unscented, and anointing breasts with scented balms or perfumes should likely be avoided, at least during the first days postpartum. However, artificial scents that mothers wear may be rapidly learnt by infants⁸¹ and appear to provide additional social cues (rather than superseding the natural breast chemosignal⁸²). What may be most important, then, is consistency of product use: habitually changing products to suit social context or mood will likely become confusing for infants.

Characterising the chemical mechanisms

Further research is required to characterise the volatile compound(s) produced by the lactating breast to which human newborns react, which would be a strong candidate for a mammary pheromone, similar to that in rabbits.¹⁹ Whether humans communicate using pheromones has been identified as one of science's outstanding questions,⁸³ and there is consensus that a breast odour signal to infants is the likeliest candidate.¹³

However, identifying a human breast chemosignal may be more challenging than in rabbits. There are no guarantees that the active signal is comprised of a single compound, as in the rabbit: it may involve two or more compounds, perhaps occurring in specific ratios (as in some other social signals⁸⁴). It may also be relatively transient, being expressed only shortly after birth or in bursts when infants are hungry. We also face additional practical and ethical constraints in research involving human newborns and parents. Thus, while a convincing characterisation of a human chemosignal would depend on a similar systematic, stepwise approach¹³ as used in the rabbit research, achieving it may require innovative methodological solutions.

One such advance could involve replacing the use of gas chromatography–mass spectrometry (currently used in most studies on mammalian olfactory communication and limited to providing single-timepoint analyses of odour samples) with continuous, real-time measurement of odour changes in undisturbed individuals, using techniques drawn from environmental and atmospheric chemistry.⁸⁵ This approach would enable the identification of compounds that are expressed at higher levels in specific relevant contexts and for short periods, such as when babies become distressed or hungry. We must also identify the appropriate methods to optimally evaluate responses to odours, which may involve working with sleeping newborns because psychophysiological measures are challenging in awake infants and stimulus-specific autonomic responses appear more reliable during sleep.⁴⁵ Finally, a conclusive identification is likely to depend on confirmation that the chemosignal is produced and responded to in the same way across genetically and culturally diverse human populations.

Although these challenges are significant, success could provide novel approaches to address breastfeeding difficulties directly. For example, the successful identification of a mammary pheromone could enable the development of a synthetic version that is attractive to newborns; this could be applied to the breasts of mothers whose babies refuse to feed or have latching difficulties, or to boost maternal self-confidence in the first few days (especially in primiparas). It might also be used to supplement the odour of mothers who are ill, who give birth prematurely or by caesarean section, who are exposed to stressful situations that interfere

with lactation,⁸⁶ or who attempt induced lactation or relactation (e.g. adoptive mothers, same-sex couples).⁸⁷

Similarly, being able to detect an infant-derived chemosignal of need might provide early warnings of hunger or distress that could, for example, be used to remotely monitor the arousal of premature babies. A synthetic version of this chemosignal could be deployed to help induce, sustain, or increase lactation and milk yield, or to assist with breast-pumping to obtain sufficient milk for a separated (e.g. preterm or ill) infant.⁸⁸

While these varied possibilities are intriguing, we would be remiss if we did not also flag potential concerns regarding undesirable applications and consequences. Foremost among these is that the development of a synthetic breast attractant should never undermine efforts to support routine breastfeeding initiation and maintenance. All aspects related to potential commercialisation should be subject to rigorous ethical scrutiny before any implementation.

Implications for international breastfeeding policy

In our view, the starkest representation of a persistent underappreciation of maternal-infant odour communication is in policy-making around breastfeeding. As described above, this underappreciation is perhaps best illustrated by the fact that there is no explicit reference to olfaction in the headline documents underpinning the WHO/UNICEF's BFHI, neither in the *Ten Steps to Successful Breastfeeding* nor in the implementation guidance.⁹

We therefore suggest that a focused analysis of policy documentation from an olfactory perspective is urgently needed. To pick one notable example, it should now be clear why we would recommend rewording of information in the section on Step 4 (immediate postnatal care) in the BFHI Implementation Guidance, where it states: 'Immediate and uninterrupted skin-to-skin contact facilitates the newborn's natural rooting reflex that helps to imprint the behaviour of *looking for the breast*' (p.17, our italics). Such wording appears to be symptomatic of an apparent underestimation or even disregard for the important role played by the sense of smell.

Indeed, we believe that a thorough and explicit consideration of the role of odour in the early stages of breastfeeding could produce evidence and insights that eventually lead to benefits on each of the *Ten Steps* (see Fig. 3). First (Step 1), recommendations for attention towards natural body odours and optimisation of exposure to these in mother-infant dyads, incorporating at least some of the points we have reviewed here, could be integrated into hospital policies. Knowledge of odour communication could be incorporated into the training of healthcare professionals to enhance staff awareness



Fig. 3: Ten ways that enhanced understanding and awareness of mother-infant odour communication can already—or potentially could—contribute to the WHO/UNICEF’s *Ten Steps to Successful Breastfeeding*.

and competency, and odour awareness training and guidance could be integrated into antenatal care for prospective parents (Steps 2 and 3). Such improved awareness would be expected to inform practices involving mammary and infant odour substrates, facilitating breastfeeding initiation, colostrum intake, and reducing early fluctuations in milk yield (Steps 4 and 5). Cumulatively, these steps could be expected to reduce the need for supplementary feeding (Step 6) and improve responsive feeding (Step 8), while awareness of the importance of odour reinforces the widely (but still not universally) recognised benefits of rooming-in, while also clarifying why rooming-in and SSC are so critical (Step 7). Furthermore, such measures should reduce the need for bottle-feeding, while maternal odour can be used, if necessary, to facilitate breast–bottle transitions (Step 9). Finally, reciprocal exposure to maternal and neonatal odours may reduce the time to discharge from hospital (Step 10), especially in premature infants (e.g. if further studies confirm that exposure accelerates the transition from tube-feeding to autonomous breastfeeding).

Conclusions and future directions

In humans, as in other mammals, olfaction is involved in the earliest interactions between mothers and infants. It helps infants overcome life’s first great behavioural challenge, in finding the nipple, plays a key

role in the subsequent establishment and maintenance of breastfeeding, and contributes to processes involved in attachment and bonding. Although further research is needed to confirm their effectiveness in clinical practice, we have outlined several ways in which understanding of maternal-infant olfactory communication could inform practical initiatives in breastfeeding initiation and maternal bonding, with corollary benefits for the health and well-being of both mothers and infants.

Further translational research awaits the discovery of the underlying chemical mechanisms. Research is thus required to characterise the volatile organic compounds produced by the lactating breast to which human newborns react, as well as those produced by infants that are reactogenic to mothers. A roadmap for such efforts has been established in the characterisation of the rabbit mammary pheromone, and successful identification could open up novel and practical ways to further facilitate olfactory mediation of breastfeeding initiation, bonding, and responsiveness of primary caretakers.

In the meantime, despite knowledge about the role of odour in nipple-finding, breastfeeding, and bonding that has accumulated over several decades, many clinicians and researchers appear to remain unaware, or at least unconvinced, of its importance, and consideration of olfaction is absent in the most recent WHO/UNICEF BFHI implementation guidance and *Ten*

Search strategy and selection criteria

Literature included in this Personal View was identified by previous research by the authors, supplemented by searches of PubMed, Web of Science, and the references of relevant articles using the search terms “odo*r”, “olfact*”, “skin-to-skin”, and “skin to skin”. We referred to papers written in languages other than English where appropriate. Abstracts and meeting reports were included where necessary to support our statements or views.

Steps. There is thus a need for greater awareness and enhanced understanding of the critical role of mammary and/or neonatal chemosignals among practitioners, parents, and policymakers alike.

Among practitioners, awareness should be increased at every stage, from hospital policies to training programs for clinical staff. Among parents and caretakers, raising awareness of early olfactory exchanges can begin during antenatal care or breastfeeding counseling and continue long after the return home, when competing or additional socio-cultural odour-related attitudes and practices may come into play. Finally, we advocate that policymakers embed an understanding and awareness of olfactory processes within the BFHI and the *Ten Steps*. By so doing, we can transform an approach to perinatal and postnatal care that currently appears to neglect this fundamental and hidden biological mechanism into one that actively leverages it, not only to promote breastfeeding rates but also to improve bonding, resilience, and the health of mothers and infants worldwide.

Outstanding questions

The outstanding question we have discussed is: how can we best apply our knowledge of olfactory communication between mothers and newborns to maximise success in breastfeeding and bonding? Important subsidiary questions include: (1) How can such knowledge be optimally incorporated into WHO/UNICEF policy, particularly the BFHI guidance and the Ten Steps? (2) What chemical compounds are responsible for the odour signal that attracts newborns to the breast immediately following birth, and can they be used to facilitate breastfeeding initiation? (3) Do infants produce an olfactory signal of need that influences maternal attentiveness?

Contributors

SCR, BS, and JH conceptualised the paper. SCR and BS wrote the original draft. SCR, BS, FD, KD, JH, DK, BL, AS, PS, VS, and JW acquired funding. FD, KD, JH, DK, BL, AS, PS, VS, JW, TA-M, DB, DD, LJ, SK, LK, ACP, KR, JCS, DS, MS, NW, and ES reviewed the draft paper and gave inputs. SCR, JH, and BS edited the final version. All authors read and approved the final version of the manuscript.

Declaration of interests

The authors declare that they have no competing interests.

Acknowledgements

The authors are funded by the Wellcome Trust (grant number: 305026/Z/23/Z). The funder had no role in study design, review, data curation and analysis, interpretation, decision to publish, or preparation of the manuscript.

References

- 1 Victora CG, Bahl R, Barros AJD, et al. Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. *Lancet*. 2016;387:475–490. [https://doi.org/10.1016/S0140-6736\(15\)01024-7](https://doi.org/10.1016/S0140-6736(15)01024-7).
- 2 UNICEF global databases. <https://data.unicef.org/topic/nutrition/breastfeeding>; 2024.
- 3 Michel MP, Gremmo-Féger G, Oger E, Sizon J. [Pilot study of early breastfeeding difficulties of term newborns: incidence and risk factors]. *Arch Pediatr*. 2007;14(5):454–460. <https://doi.org/10.1016/j.arcped.2007.01.005>.
- 4 Dewey KG, Nommsen-Rivers LA, Heinig MJ, Cohen RJ. Risk factors for suboptimal infant breastfeeding behavior, delayed onset of lactation, and excess neonatal weight loss. *Pediatrics*. 2003;112(3 Pt 1):607–619. <https://doi.org/10.1542/peds.112.3.607>.
- 5 Aderibigbe T, Srisopa P, Henderson WA, Lucas R. Meta-ethnography on the experiences of women from around the world who exclusively breastfed their full-term infants. *J Obstet Gynecol Neonatal Nurs*. 2024;53(2):120–131. <https://doi.org/10.1016/j.jogn.2023.11.008>.
- 6 Vögele J, Rittershaus L, Halling T. “Breast is best”—Infant-feeding, infant mortality and infant welfare in Germany during the late nineteenth and twentieth centuries. *Health*. 2013;5:2190–2203.
- 7 Schaal B, Al Ain S. Chemical signals ‘selected for’ newborns in mammals. *Anim Behav*. 2014;97:289–299. <https://doi.org/10.1016/j.anbehav.2014.08.022>.
- 8 Schaal B, Saxton TK, Loos H, Soussignan R, Durand K. Olfaction scaffolds the developing human from neonate to adolescent and beyond. *Philos Trans R Soc Lond B Biol Sci*. 2020;375(1800):20190261. <https://doi.org/10.1098/rstb.2019.0261>.
- 9 WHO/UNICEF. *Protecting, promoting and supporting breastfeeding in facilities providing maternity and newborn services: implementing the revised Baby-friendly hospital initiative*. 2018.
- 10 McGann JP. Poor human olfaction is a 19th-century myth. *Science*. 2017;356(6338):eaam7263.
- 11 Bratman GN, Bembibre C, Daily GC, et al. Nature and human well-being: the olfactory pathway. *Sci Adv*. 2024;10(20):eadn3028. <https://doi.org/10.1126/sciadv.adn3028>.
- 12 Loos HM, Schaal B, Pause BM, et al. Past, present, and future of human chemical communication research. *Perspect Psychol Sci*. 2025;20(1):20–44. <https://doi.org/10.1177/17456916231188147>.
- 13 Wyatt TD. The search for human pheromones: the lost decades and the necessity of returning to first principles. *Proc Biol Sci*. 2015;282(1804):20142994. <https://doi.org/10.1098/rspb.2014.2994>.
- 14 Roberts SC, Havlíček J, Schaal B. Human olfactory communication: current challenges and future prospects. *Philos Trans R Soc Lond B Biol Sci*. 2020;375(1800):20190258. <https://doi.org/10.1098/rstb.2019.0258>.
- 15 Schneider NY, Fletcher TP, Shaw G, Renfree MB. The olfactory system of the tamar wallaby is developed at birth and directs the neonate to its mother’s pouch odours. *Reproduction*. 2009;138(5):849–857. <https://doi.org/10.1530/REP-09-0145>.
- 16 Galef BG, Muskus PA. Olfactory mediation of mother–young contact in long-evans rats. *J Comp Physiol Psychol*. 1979;93:708–716.
- 17 Teicher MH, Flaum LE, Williams M, Eckert SJ, Lumia AR. Survival, growth and suckling behavior of neonatal bulbectomized rats. *Physiol Behav*. 1978;21(4):553–561. [https://doi.org/10.1016/0031-9384\(78\)90130-0](https://doi.org/10.1016/0031-9384(78)90130-0).
- 18 Blass EM, Shuleikina-Turpaeva K, Lusckekin V. Sensory determinants of nipple-attachment behavior in 2–4-day-old kittens. *Dev Psychobiol*. 1988;21(4):365–370. <https://doi.org/10.1002/dev.420210407>.
- 19 Schaal B, Coureaud G, Langlois D, Giniès C, Sémon E, Perrier G. Chemical and behavioural characterisation of the rabbit mammary pheromone. *Nature*. 2003;424:68–72.

- 20 Coureaud G, Langlois D, Perrier G, Schaal B. Convergent changes in the maternal emission and pup reception of the rabbit mammary pheromone. *Chemoecology*. 2006;16(3):169–174. <https://doi.org/10.1007/s00049-006-0345-9>.
- 21 Coureaud G, Fortun-Lamothe L, Langlois D, Schaal B. The reactivity of neonatal rabbits to the mammary pheromone as a probe for viability. *Animal*. 2007;1(7):1026–1032. <https://doi.org/10.1017/s1751731107000389>.
- 22 Coureaud G, Schaal B, Langlois D, Perrier G. Orientation response of newborn rabbits to odours of lactating females: relative effectiveness of surface and milk cues. *Anim Behav*. 2001;61(1):153–162. <https://doi.org/10.1006/anbe.2000.1563>.
- 23 Schaal B. Mammary odor cues and pheromones: mammalian infant-directed communication about maternal state, mammae, and milk. *Vitam Horm*. 2010;83:83–136. [https://doi.org/10.1016/S0083-6729\(10\)83004-3](https://doi.org/10.1016/S0083-6729(10)83004-3).
- 24 Marlier L, Schaal B, Soussignan R. Bottle-fed neonates prefer an odor experienced in utero to an odor experienced postnatally in the feeding context. *Dev Psychobiol*. 1998;33(2):133–145.
- 25 Soussignan R, Schaal B, Marlier L, Jiang T. Facial and autonomic responses to biological and artificial olfactory stimuli in human neonates: re-examining early hedonic discrimination of odors. *Physiol Behav*. 1997;62(4):745–758. [https://doi.org/10.1016/s0031-9384\(97\)00187-x](https://doi.org/10.1016/s0031-9384(97)00187-x).
- 26 Doucet S, Soussignan R, Sagot P, Schaal B. The "smellscape" of mother's breast: effects of odor masking and selective unmasking on neonatal arousal, oral, and visual responses. *Dev Psychobiol*. 2007;49(2):129–138. <https://doi.org/10.1002/dev.20210>.
- 27 Righard L, Alade MO. Effect of delivery room routines on success of first breast-feed. *Lancet*. 1990;336(8723):1105–1107. [https://doi.org/10.1016/0140-6736\(90\)92579-7](https://doi.org/10.1016/0140-6736(90)92579-7).
- 28 Widström AM, Lilja G, Aaltomaa-Michalios P, Dahllöf A, Lintula M, Nissen E. Newborn behaviour to locate the breast when skin-to-skin: a possible method for enabling early self-regulation. *Acta Paediatr*. 2011;100(1):79–85. <https://doi.org/10.1111/j.1651-2227.2010.01983.x>.
- 29 Varendi H, Porter RH, Winberg J. Does the newborn baby find the nipple by smell? *Lancet*. 1994;344(8928):989–990. [https://doi.org/10.1016/s0140-6736\(94\)91645-4](https://doi.org/10.1016/s0140-6736(94)91645-4).
- 30 Varendi H, Porter RH. Breast odour as the only maternal stimulus elicits crawling towards the odour source. *Acta Paediatr*. 2001;90(4):372–375.
- 31 Hym C, Forma V, Anderson DI, et al. Newborn crawling and rooting in response to maternal breast odor. *Dev Sci*. 2021;24(3):e13061. <https://doi.org/10.1111/desc.13061>.
- 32 Baudesson de Chanville A, Brevaut-Malaty V, Garbi A, Tosello B, Baumstarck K, Gire C. Analgesic effect of maternal human milk odor on premature neonates: a randomized controlled trial. *J Hum Lact*. 2017;33(2):300–308. <https://doi.org/10.1177/0890334417693225>.
- 33 Schaal B, Marlier L, Soussignan R. Olfactory function in the human fetus: evidence from selective neonatal responsiveness to the odor of amniotic fluid. *Behav Neurosci*. 1998;112(6):1438–1449. <https://doi.org/10.1037/0735-7044.112.6.1438>.
- 34 Sullivan RM, Taborsky-Barba S, Mendoza R, et al. Olfactory classical conditioning in neonates. *Pediatrics*. 1991;87(4):511–518. [https://doi.org/10.1016/s0149-7634\(98\)00044-x](https://doi.org/10.1016/s0149-7634(98)00044-x).
- 35 Porter RH, Winberg J. Unique salience of maternal breast odors for newborn infants. *Neurosci Biobehav Rev*. 1999;23(3):439–449. [https://doi.org/10.1016/s0149-7634\(98\)00044-x](https://doi.org/10.1016/s0149-7634(98)00044-x).
- 36 Schaal B. Pheromones for newborns. In: Mucignat-Caretta C, ed. *Neurobiology of chemical communication*. Boca Raton: CRC Press; 2014:483–515.
- 37 Marlier L, Schaal B. Human newborns prefer human milk: conspecific milk odor is attractive without postnatal exposure. *Child Dev*. 2005;76(1):155–168. <https://doi.org/10.1111/j.1467-8624.2005.00836.x>.
- 38 Makin JW, Porter RH. Attractiveness of lactating females' breast odors to neonates. *Child Dev*. 1989;60(4):803–810.
- 39 Mizuno K, Ueda A. Antenatal olfactory learning influences infant feeding. *Early Hum Dev*. 2004;76(2):83–90. <https://doi.org/10.1016/j.earlhumdev.2003.10.003>.
- 40 Loos HM, Reger D, Schaal B. The odour of human milk: its chemical variability and detection by newborns. *Physiol Behav*. 2019;199:88–99. <https://doi.org/10.1016/j.physbeh.2018.11.008>.
- 41 Oftedal OT, Iverson SJ. Comparative analysis of nonhuman milks: a phylogenetic variation in gross composition of milks. In: Jensen RG, ed. *Handbook of Milk Composition*. Academic Press; 1995:749–788.
- 42 Loos H. *Ethological and aroma-analytical investigations on human milk odour*. PhD thesis. Friedrich-Alexander University; 2015.
- 43 Klaey-Tassone M, Durand K, Damon F, et al. Human neonates prefer colostrum to mature milk: evidence for an olfactory bias toward the "initial milk"? *Am J Hum Biol*. 2021;33(5):e23521. <https://doi.org/10.1002/ajhb.23521>.
- 44 Schaal B, Doucet S, Sagot P, Hertling E, Soussignan R. Human breast areolae as scent organs: morphological data and possible involvement in maternal-neonatal coadaptation. *Dev Psychobiol*. 2006;48(2):100–110. <https://doi.org/10.1002/dev.20122>.
- 45 Doucet S, Soussignan R, Sagot P, Schaal B. The secretion of areolar (Montgomery's) glands from lactating women elicits selective, unconditional responses in neonates. *PLoS One*. 2009;4(10):e7579. <https://doi.org/10.1371/journal.pone.0007579>.
- 46 Poindron P, Lévy F, Krehbiel D. Genital, olfactory, and endocrine interactions in the development of maternal behaviour in the parturient ewe. *Psychoneuroendocrinology*. 1988;13(1–2):99–125. [https://doi.org/10.1016/0306-4530\(88\)90009-1](https://doi.org/10.1016/0306-4530(88)90009-1).
- 47 Brouette-Lahlou I, Amouroux R, Chastrette F, Cosnier J, Stoffelsma J, Vernet-Maury E. Dodecyl propionate, attractant from rat pup preputial gland: characterization and identification. *J Chem Ecol*. 1991;17(7):1343–1354. <https://doi.org/10.1007/BF00983767>.
- 48 Croy I, Mohr T, Weidner K, Hummel T, Junge-Hoffmeister J. Mother-child bonding is associated with the maternal perception of the child's body odor. *Physiol Behav*. 2019;198:151–157. <https://doi.org/10.1016/j.physbeh.2018.09.014>.
- 49 Lundström JN, Mathe A, Schaal B, et al. Maternal status regulates cortical responses to the body odor of newborns. *Front Psychol*. 2013;4:597. <https://doi.org/10.3389/fpsyg.2013.00597>.
- 50 Schäfer L, Köppel C, Krefner-Kiel D, et al. The scent of cuteness—neural signatures of infant body odors. *Soc Cogn Affect Neurosci*. 2024;19(1):nsae038. <https://doi.org/10.1093/scan/nsae038>.
- 51 Dubas JS, Heijkoop M, van Aken MAG. A preliminary investigation of parent-progeny olfactory recognition and parental investment. *Hum Nat*. 2009;20(1):80–92. <https://doi.org/10.1007/s12110-009-9058-4>.
- 52 Schneeberger K, Röder G, Taborsky M. The smell of hunger: Norway rats provision social partners based on odour cues of need. *PLoS Biol*. 2020;18(3):e3000628. <https://doi.org/10.1371/journal.pbio.3000628>.
- 53 Pabio BM, Schäfer AS, Hoenen M, Lübke KT, Stockhorst U. You see what you smell: preferential processing of chemosensory satiety cues and its impact on body shape perception. *Brain Sci*. 2021;11(9):1152. <https://doi.org/10.3390/brainsci11091152>.
- 54 Roberts SC, Trěbická Fialová J, Sorokowska A, et al. Emotional expression in human odour. *Evol Hum Sci*. 2022;4:e44. <https://doi.org/10.1017/ehs.2022.44>.
- 55 Lübke KT, Busch A, Hoenen M, Schaal B, Pause BM. Chemosensory anxiety signals prime defensive behavior in prepubertal girls. *Physiol Behav*. 2017;173:30–33. <https://doi.org/10.1016/j.physbeh.2017.01.035>.
- 56 Abdulghani N, Edvardsson K, Amir LH. Worldwide prevalence of mother-infant skin-to-skin contact after vaginal birth: a systematic review. *PLoS One*. 2018;13(10):e0205696. <https://doi.org/10.1371/journal.pone.0205696>.
- 57 Packheiser J, Hartmann H, Fredriksen K, Gazzola V, Keyser C, Michon F. A systematic review and multivariate meta-analysis of the physical and mental health benefits of touch interventions. *Nat Hum Behav*. 2024;8(6):1088–1107. <https://doi.org/10.1038/s41562-024-01841-8>.
- 58 Moberg KU, Handlin L, Petersson M. Neuroendocrine mechanisms involved in the physiological effects caused by skin-to-skin contact - with a particular focus on the oxytocinergic system. *Infant Behav Dev*. 2020;61:101482. <https://doi.org/10.1016/j.infbeh.2020.101482>.
- 59 Moore ER, Bergman N, Anderson GC, Medley N. Early skin-to-skin contact for mothers and their healthy newborn infants. *Cochrane Database Syst Rev*. 2016;11(11):Cd003519. <https://doi.org/10.1002/14651858.CD003519.pub4>.
- 60 Widström AM, Brimdyr K, Svensson K, Cadwell K, Nissen E. Skin-to-skin contact the first hour after birth, underlying implications and clinical practice. *Acta Paediatr*. 2019;108(7):1192–1204. <https://doi.org/10.1111/apa.14754>.
- 61 Klaus M. Mother and infant: early emotional ties. *Pediatrics*. 1998;102(5 Suppl E):1244–1246.
- 62 Rana P, Swain D. Efficacy of the standard breast crawl technique on maternal and newborn outcomes after term vaginal birth: a

- randomized controlled trial. *J Midwifery Womens Health*. 2023;68(4):473–479. <https://doi.org/10.1111/jmwh.13487>.
- 63 Klaus MH, Jerauld R, Kreger NC, McAlpine W, Steffa M, Kennel JH. Maternal attachment. Importance of the first postpartum days. *N Engl J Med*. 1972;286(9):460–463. <https://doi.org/10.1056/nejm197203022860904>.
- 64 Bystrova K, Ivanova V, Edhborg M, et al. Early contact versus separation: effects on mother-infant interaction one year later. *Birth*. 2009;36(2):97–109. <https://doi.org/10.1111/j.1523-536X.2009.00307.x>.
- 65 Mizuno K, Mizuno N, Shinohara T, Noda M. Mother-infant skin-to-skin contact after delivery results in early recognition of own mother's milk odour. *Acta Paediatr*. 2004;93(12):1640–1645. <https://doi.org/10.1080/08035250410023115>.
- 66 Bartocci M, Winberg J, Ruggiero C, Bergqvist LL, Serra G, Lagercrantz H. Activation of olfactory cortex in newborn infants after odor stimulation: a functional near-infrared spectroscopy study. *Pediatr Res*. 2000;48(1):18–23. <https://doi.org/10.1203/00006450-200007000-00006>.
- 67 Welch MG, Hofer MA, Brunelli SA, et al. Family nurture intervention (FNI): methods and treatment protocol of a randomized controlled trial in the NICU. *BMC Pediatr*. 2012;12:14. <https://doi.org/10.1186/1471-2431-12-14>.
- 68 Beebe B, Myers MM, Lee SH, et al. Family nurture intervention for preterm infants facilitates positive mother-infant face-to-face engagement at 4 months. *Dev Psychol*. 2018;54(11):2016–2031. <https://doi.org/10.1037/dev0000557>.
- 69 Woodward LJ, Ludwig RJ, Gong A, Myers MM, Welch MG. Family Nurture Intervention and mother-preterm infant autonomic emotional connection in the neonatal intensive care unit: a review. *Acta Paediatr*. 2025;114(4):760–771. <https://doi.org/10.1111/apa.17456>.
- 70 Welch MG, Firestein MR, Austin J, et al. Family Nurture Intervention in the Neonatal Intensive Care Unit improves social-relatedness, attention, and neurodevelopment of preterm infants at 18 months in a randomized controlled trial. *J Child Psychol Psychiatry*. 2015;56(11):1202–1211. <https://doi.org/10.1111/jcpp.12405>.
- 71 Letzkus L, Conaway M, Miller-Davis C, Darring J, Keim-Malpass J, Zanelli S. A feasibility randomized controlled trial of a NICU rehabilitation program for very low birth weight infants. *Sci Rep*. 2022;12(1):1729. <https://doi.org/10.1038/s41598-022-05849-w>.
- 72 Qin Y, Liu S, Yang Y, Zhong Y, Hao D, Han H. Effects of human milk odor stimulation on feeding in premature infants: a systematic review and meta-analysis. *Sci Rep*. 2024;14(1):8964. <https://doi.org/10.1038/s41598-024-59175-4>.
- 73 Delgado Paramo L, Bronnert A, Lin L, Bloomfield FH, Muelbert M, Harding JE. Exposure to the smell and taste of milk to accelerate feeding in preterm infants. *Cochrane Database Syst Rev*. 2024;5(5):Cd013038. <https://doi.org/10.1002/14651858.CD013038.pub3>.
- 74 Das S, Bloomfield FH, Alexander T, et al. Effect of different feeding strategies on neurosensory impairment at 2 years' corrected age in children born moderate-and-late preterm: follow-up of the DIAMOND randomised trial. *J Paed Child Health*. 2025;61(S1):11.
- 75 Zhang S, Su F, Li J, Chen W. The analgesic effects of maternal milk odor on newborns: a meta-analysis. *Breastfeed Med*. 2018;13(5):327–334. <https://doi.org/10.1089/bfm.2017.0226>.
- 76 Jessen S. Maternal odor reduces the neural response to fearful faces in human infants. *Dev Cogn Neurosci*. 2020;45:100858. <https://doi.org/10.1016/j.dcn.2020.100858>.
- 77 Badiie Z, Asghari M, Mohammadizadeh M. The calming effect of maternal breast milk odor on premature infants. *Pediatr Neonatol*. 2013;54(5):322–325. <https://doi.org/10.1016/j.pedneo.2013.04.004>.
- 78 Kido M, Yonezawa K, Haruna M, Tahara-Sasagawa E, Usui Y. A global survey on national standard care for newborn bathing. *Jpn J Nurs Sci*. 2024;21(1):e12558. <https://doi.org/10.1111/jjns.12558>.
- 79 Brimdyr K, Cadwell K, Stevens J, Takahashi Y. An implementation algorithm to improve skin-to-skin practice in the first hour after birth. *Matern Child Nutr*. 2018;14(2):e12571. <https://doi.org/10.1111/mcn.12571>.
- 80 Kaya A, Yeşildere Sağlam H, Karadağ E, Gürsoy E. The effectiveness of aromatherapy in the management of labor pain: a meta-analysis. *Eur J Obstet Gynecol Reprod Biol X*. 2023;20:100255. <https://doi.org/10.1016/j.eurox.2023.100255>.
- 81 Schleidt M, Genzel C. The significance of mother's perfume for infants in the first weeks of their life. *Ethol Sociobiol*. 1990;11(3):145–154. [https://doi.org/10.1016/0162-3095\(90\)90007-5](https://doi.org/10.1016/0162-3095(90)90007-5).
- 82 Delaunay-El Allam M, Marlier L, Schaal B. Learning at the breast: preference formation for an artificial scent and its attraction against the odor of maternal milk. *Infant Behav Dev*. 2006;29(3):308–321. <https://doi.org/10.1016/j.infbeh.2005.12.008>.
- 83 Seife C, Anon. So much more to know. *Science*. 2005;309:78–102.
- 84 Hölldobler B. The chemistry of social regulation: multicomponent signals in ant societies. *Proc Natl Acad Sci U S A*. 1995;92(1):19–22. <https://doi.org/10.1073/pnas.92.1.19>.
- 85 Roberts SC, Misztal PK, Langford B. Decoding the social volatome by tracking rapid context-dependent odour change. *Philos Trans R Soc Lond B Biol Sci*. 2020;375(1800):20190259. <https://doi.org/10.1098/rstb.2019.0259>.
- 86 Erat Nergiz M, Barutçu A, Güneş B, et al. Breastfeeding problems and associated factors: a cross-sectional study after the 2023 Türkiye earthquake. *Int Breastfeed J*. 2025;20(1):32. <https://doi.org/10.1186/s13006-025-00724-6>.
- 87 Lawrence RA. Induced lactation and relactation (including nursing an adopted baby) and cross-nursing. In: Lawrence RA, Lawrence RM, eds. *Breastfeeding*. 9th ed. Elsevier; 2022:628–645.
- 88 Hoban R, Medina-Poeliniz C, Signorile M, Janes J, Fan C-PS, Meier PP. Early postpartum pumping behaviors, pumped milk volume, and achievement of secretory activation in breast pump-dependent mothers of preterm infants. *J Perinatol*. 2024;44(11):1597–1606. <https://doi.org/10.1038/s41372-024-02021-2>.