

Wetland landscape transformation by beavers: responses of biodiversity and functional indicators at multiple scales

Received: 17 Apr 2025

Accepted: 24 Jan 2026

Alan Law, Nigel Willby, Tom Spencer, David Bryan, Garth Foster, Lori Handley, Wenfei Liao, Graham Sellers & Petri Nummi

Cite this article as: Law, A., Willby, N., Spencer, T. *et al.* Wetland landscape transformation by beavers: responses of biodiversity and functional indicators at multiple scales. *Landsc Ecol* (2026). <https://doi.org/10.1007/s10980-026-02303-4>

We are providing an unedited version of this manuscript to give early access to its findings. Before final publication, the manuscript will undergo further editing. Please note there may be errors present which affect the content, and all legal disclaimers apply.

If this paper is publishing under a Transparent Peer Review model then Peer Review reports will publish with the final article.

Title page**Title**

Wetland landscape transformation by beavers: responses of biodiversity and functional indicators at multiple scales

Author names and affiliations

Alan Law^{1*}, Nigel Willby¹, Tom Spencer³, David Bryan¹, Garth N. Foster², Lori Lawson

Handley⁶, Wenfei Liao⁴, Graham S. Sellers³ and Petri Nummi⁵

1. Biological and Environmental Sciences, University of Stirling, Stirling, FK9 4LA, United Kingdom
2. Aquatic Coleoptera Conservation Trust, Ayr, KA7 1JJ, United Kingdom
3. School of Environmental and Life Sciences, University of Hull, Hull, HU6 7RX, United Kingdom
4. Department of Geosciences and Geography, University of Helsinki, P.O.Box 64, FI-00014, Finland
5. Department of Forest Sciences, University of Helsinki, P.O.Box 27, FI-00014, Finland
6. Aquatic Ecosystems Group, UK Centre for Ecology & Hydrology, Library Avenue, Bailrigg, Lancaster, LA1 4AP, United Kingdom.

* Corresponding author: Alan Law, Biological and Environmental Sciences, Cottrell building, University of Stirling, Stirling, FK9 4LA, UK. (0044) 01786 467 770, alan.law1@stir.ac.uk. ORCID ID: [0000-0001-5971-3214](https://orcid.org/0000-0001-5971-3214)

Abstract

Context

Landscape-scale restoration is needed to reverse declines in biodiversity, but the ecological processes that sustain biodiversity by boosting heterogeneity are often overlooked. Large herbivores are important drivers of heterogeneity and are increasingly being used to restore lost dynamic processes.

Objectives

With beaver populations recovering from a historic low, we test what their ecosystem engineering potential means for biodiversity and ecosystem functioning at multiple scales.

Methods

We quantified 10 taxonomic groups at sample, site and landscape scale via in-situ surveys (plants and water beetles) and eDNA sampling (invertebrate and vertebrates) from nine beaver-created wetlands and nine wetlands unmodified by beavers (control wetlands) in Evo, Finland.

Results

Per taxonomic group, the mean and total number of taxa at sample and site-scale was mostly similar between wetland types, though significantly higher in beaver wetlands at sample (true flies) and site-scale (plants and true flies). 63% of all taxa were shared by beaver-created and control wetlands. However, both wetland types supported unique taxa with beaver wetlands increasing the landscape taxon pool by an average of 19% (range 0-

40%), most notably for plants, beetles, true flies and may/stone/caddisflies. Plant functional diversity was 55% higher in beaver compared to control wetlands.

Conclusions

Beaver wetlands are integral to reinstating dynamic ecological processes that provide refugia for multiple taxonomic groups, supporting taxa otherwise absent from the landscape. Our findings hint at the scale of past biodiversity loss associated with beaver-dependent wetlands, while offering a glimpse of what could be gained from their ongoing population recovery.

Keywords

Boreal wetlands, community composition, ecosystem engineer, biodiversity, functional diversity, restoration

Main text

Introduction

The recovery of freshwater biodiversity and ecosystem functioning is arguably the most fundamental challenge in restoration ecology (Tickner et al. 2020; Maasri et al. 2022). Freshwaters are disproportionately biodiverse environments, providing critical resources (Dudgeon 2019), yet their freshwater biodiversity is declining twice as fast compared to terrestrial and marine environments (Leung et al. 2020). Protecting existing ecosystems, habitats and species is insufficient to reverse biodiversity declines, therefore ecosystem restoration is necessary (Young 2000; Mutillod et al. 2024). Ecosystem restoration principles are broadly well known, such as peatland rewetting, native tree regeneration, restoring river floodplain connectivity, and biomanipulating shallow lake trophic cascades. These principles are usually supported by habitat-specific methods and guidance, although outcomes are often variable (Palmer et al. 2014; Sinclair et al. 2023). Demonstrating wider effects of restoration beyond the focal habitat or species—for example, the re-establishment of functional processes (disturbance, herbivory, predation) that sustain ecosystems—is beyond the success criteria or monitoring and reporting scope of most restoration projects. However, these processes are essential if ecosystem restoration is to become self-sustaining and to progress at the scale needed.

Whilst the concept of ecosystem restoration is not new, the growing use of ecosystem engineers as nature-based solutions or as an active part of rewilding has prompted new perspectives and stimulated interest in restorative processes (Byers et al. 2006; Mutillod et al. 2024). Particularly in countries where such species have been lost and the chances of natural comeback are low (Gaywood and Stanley-Price 2022). This has

created an impetus where numerous ecological restoration or rewilding projects are underway or proposed, designed to re-energise natural processes to improve the functionality of ecosystems. Various large herbivores, including moose, bison and boar, or their domesticated functional analogues, act as agents for these processes, although beavers are widely used, having now been reintroduced or spread naturally to ~28 countries across Europe (Rosell and Campbell-Palmer 2022). Coupled with their capacity to engineer ecosystems via wetland creation, the recovery of beaver populations has consequences for ecosystem processes spanning herbivory, decomposition, nutrient cycling and succession (Larsen et al. 2021). However, with many such ecological restoration projects being less than 10 years old and therefore still in their infancy, the need to understand the longer-term implications for biodiversity of renewed ecological processes in modern landscapes is becoming timely. Particularly as rewilding gains momentum and projects are distributed across a widening diversity of land cover types.

Beavers are the model ecosystem-engineer, second only to humans in their capacity to transform landscapes and increasingly proclaimed as part of the solution to the freshwater biodiversity crisis (Law et al. 2019; Orazi et al. 2022; Andersen et al. 2023). Beavers engineer landscapes by impounding streams with woody dams thereby creating shallow wetlands. Dam maintenance, canal digging, tree felling and selective foraging create further disturbances at both small- and large-scales contributing to a unique wetland environment with strong, positive effects on freshwater biodiversity (Stringer and Gaywood 2016; Law et al. 2017; Willby et al. 2018; Nummi and Holopainen 2020). As the effects of ecosystem engineering by beavers on multiple ecosystem processes are increasingly recognised (Byers 2022), reintroduction projects have moved beyond a focus on population

recovery, to positioning beavers as the architects of dynamic, resilient environments in restoration and rewilding projects across Europe and North America (Burchsted et al. 2010).

The physical changes to landscapes following beaver occupation are well documented, with dams and wetlands stabilising flows (Puttock et al. 2020), increasing sediment storage (Butler and Malanson 1995; Puttock et al. 2018), altering geomorphology, hydrology and biogeochemistry (Brazier et al. 2020; Larsen et al. 2021), and restructuring aquatic and riparian vegetation (Wright et al. 2002; Nummi and Kuuluvainen 2013; Wilson et al. 2024) to create highly heterogenous environments. The role of beavers in wetland creation and habitat disturbance has often been gauged via biodiversity responses. Changes to biota in the riparian (McCaffery and Eby 2016; Wikar et al. 2024), terrestrial (Orazi et al. 2022; Fedyń et al. 2023), arboreal (Harper et al. 2019b) and aerial (Nummi et al. 2011; Ciechanowski et al. 2011; Moser et al. 2025) zones have all been well documented. Studies within beaver wetlands have assessed changes in vegetation over time (Ray et al. 2001; Law et al. 2017) or space by comparing multiple beaver-created/altered habitats to non-beaver influenced sites (Hood & Larson, 2013; Nummi & Holopainen 2014; Bush & Wissinger, 2016; Law, McLean & Willby, 2016; Washko, Roper & Atwood, 2020). More focussed studies also report the use of beaver dams, lodges and feeding debris (e.g. dead wood) by other species (Thompson et al. 2016; Schloemer et al. 2023; Wilson and Bremner-Harrison 2024; Andersen et al. 2024). Within these examples, the focal species range through plants, invertebrates, amphibians, fish, birds and mammals, with most studies focussing on responses within individual sites or single taxonomic groups, the only notable exception being on terrestrial biodiversity (Orazi et al. 2022). Fewer studies compare different elements of biodiversity among beaver wetlands in the same (Nummi and Holopainen 2014; Nummi et al. 2021b; Spyra et al. 2023) or different landscapes (Nummi et al. 2019b), and

fewer still consider the comparative biodiversity of beaver wetlands and similar wetland types formed independently of beavers in the same landscape (Willby et al. 2018; Law et al. 2019; Nummi et al. 2019a). Consequently, effects on disparate biota in parallel are poorly understood, including for mobile taxa that interact more broadly with beaver wetlands (i.e. birds, mammals) at a landscape scale (Fedyń et al. 2024; Kivinen and Nummi 2025).

Most studies on ecosystem responses to beaver engineering have also focussed on biodiversity from a taxonomic perspective (e.g. species richness) rarely considering the functional role and ecological preferences of species. Some have reported changes in freshwater macroinvertebrate functional feeding groups due to beaver damming (see Washko, Willby & Law (2022) for a review), but knowledge gaps remain. Functional diversity is a meaningful indicator of restoration success (Coccia et al. 2021; Lu et al. 2023), but we currently lack evidence that beaver-dependent wetlands perform a different functional role in the landscape to existing waterbodies, or if they might create more functional resilience via redundancy. This incomplete picture of the restoration potential of a key ecosystem engineer is important in the context of the drive towards landscape restoration for multiple processes and benefits (UNESCO World Water Assessment Programme 2018). Current environmental paradigms emphasise the benefits of more biodiverse, connected and complex landscapes, particularly via ecosystem restoration (e.g. UN decade on Ecosystem Restoration). Set alongside a global reduction in wetland quantity and quality (Gardner and Finlayson 2018), beaver-dependent wetlands might therefore enhance landscape resilience by acting as highly connected and biodiverse hotspots.

Using in situ surveys (aquatic plants, beetles) and eDNA sampling (vertebrates and invertebrates), the primary aim of our study was to quantify differences in biodiversity from across multiple species groups from beaver-created and control wetlands at sample-, site-

and landscape-scale. Differences in functional diversity were inferred from ecological traits from plants only. We hypothesised that; i) both wetland types (beaver and control) will contain unique species ii) creation legacy and ongoing disturbance will facilitate distinct biodiversity and plant function in beaver wetlands, and iii) the presence of beaver-created wetlands will increase landscape-scale biodiversity and function.

Methods

Study area

All data were gathered during August 2022 in the Evo catchment (area 39 km²) in southern Finland (61°12' N, 25°07' E) (Fig. 1). This area contains a mosaic of wetlands, including ~100 lakes and ponds, some beaver-created or beaver-modified. The majority are not hydrologically influenced by beavers but nested within the same landscape (Kivinen et al. 2020). The area has a cool continental climate with a mean annual temperature of 4.7°C and a mean annual precipitation of 639 mm (<https://www.ilmatieteenlaitos.fi/1991-2020-lampotilatilastot>). Waterbodies are ice-covered from November to April in line with duration of snow cover (Nummi et al. 2019a).

The lakes of the area are meso- and oligotrophic, relatively small (0.1 – 49.5 ha), and mostly hydrologically inter-connected (Nummi and Pöysä 1993; Arvola et al. 2025). Most have one or more inflows and one outflow and can be considered as surface runoff or groundwater lakes. Groundwater lakes are typically situated on the glacio-fluvial sandy deposits in the lower part of the area and surface runoff lakes on the till deposits in the upper areas (Arvola et al. 2025). Boreal forest (dominated by *Picea abies* and *Pinus*

sylvestris) covers most of the area, interspersed with mires and lakes. The influences of agriculture and human settlement are limited and local, with forestry the only major human disturbance in the area (Arvola et al. 2025). Apart from beaver-created variability, the lakes of Evo are fairly stable (Suhonen, Nummi & Pöysä, 2011; Nummi, Arzel & Sauramo, 2021b). At Evo, beavers most often create wetlands by damming existing lakes, but will also dam streams (Hyvönen and Nummi 2008; Nummi and Hahtola 2008). Within the Evo landscape, beavers typically move from one lake to another every three years on average, abandoning old sites and creating new ones or reworking former beaver habitats (Hyvönen and Nummi 2008; Kivinen et al. 2020). Within the Evo area, beavers are the North American species, *Castor canadensis*, introduced in the 1930's before the genetic separation from the native Eurasian *C. fiber* had been resolved. Only minor differences in life history, ecology and behaviour are known between these species, which suggest a comparable niche overlap and therefore functional equivalence (Parker et al. 2012).

In general, across the Evo study region the non-beaver lake and pond shores are steep, with little emergent vegetation. The narrow vegetation belts consist mainly of sedges (*Carex* spp.) and the common reed (*Phragmites australis*). Emergent vegetation can be fringed by a narrow belt of yellow (*Nuphar lutea*) and/or white water lilies (*Nymphaea candida*) but submerged vegetation is sparse. The riparian zone is typically dominated by *Sphagnum* mosses or dwarf shrubs. Beaver wetlands have significantly shallower shores (mean = 23 cm) than non-beaver wetlands (mean = 64 cm) and often contain inundated bushes and herbaceous vegetation (Nummi and Hahtola 2008). Sedges also dominate in damp meadows abandoned by beavers. All waterbodies are usually edged with a narrow belt of deciduous trees i.e. birch (*Betula* spp.), alder (*Alnus* spp.) and willow (*Salix* spp.).

Physico-chemical conditions in the control and beaver wetlands are broadly similar, though DOC and P concentrations tend to be higher in beaver wetlands (Vehkaoja et al. 2015).

Sampling design

Nine beaver-created and nine non-beaver control wetlands were surveyed (Fig. 1). Beaver wetlands comprised modified pre-existing small lakes with dammed outflows which flooded low-lying surrounding forest (Fig. 2a and c). Beaver wetlands were surveyed when considered active i.e. exhibited signs of fresh feeding, tree felling and dam maintenance. Their ages ranged from 3-35 years, with older sites having been abandoned and reflooded several times (Kivinen et al. 2020). Control wetlands were selected based on comparatively similar size, being located within the same landscape and drainage basin as the beaver wetlands, but with their hydrology unaltered by beavers (Fig. 2b and d). These represent the most closely comparable reference systems that naturally exist in this landscape independently of beavers and are therefore a suitable benchmark against which to measure any additional biodiversity benefits that arise from beavers. Further details of beaver and control sites in this region can be found in (Nummi et al. 2019a).




Figure 1. The location of the study area Evo, Finland. Beaver wetlands are denoted by a beaver symbol and control wetlands by a vegetation symbol. Numbers refer to each wetland sampled (Table S1). Contains material from the National Land Survey of Finland's Terrain Database 12/2024.




Figure 2. Examples of the studied wetlands from Evo, Finland. Aerial pictures of (A) beaver wetland Peikkoplotti (site 9), and (B) control wetland Likojärvi (site 14). Picture from the edge of (C) beaver wetland Peikkoplotti (site 9), and (D) Alin Mustajärvi (sites 8) © Alan Law.

Field methods

To determine water quality, a 500 ml water sample was collected from each wetland using a throw bottle from the shore. This was chilled immediately and analysed within 12 hrs for pH, alkalinity (mmol/L), total nitrogen ($\mu\text{g/L}$), total phosphorus ($\mu\text{g/L}$), chlorophyll A ($\mu\text{g/L}$) and dissolved organic carbon (mg/L) in the laboratory. Contextual information on tree composition and density, water depth (cm), wetland area (ha), wetland age, altitude (m

a.s.l), distance to nearest wetland (m), inflow and outflow were available from previous studies (Kivinen et al. 2020; Nummi and Holopainen 2020).

Both aquatic plants and water beetles were sampled via in-situ surveys. Both groups are diverse, their communities partly indicative of local conditions, and provide a contrast between passive and active dispersers. Each wetland was visually subdivided into dominant mesohabitats according to vegetation (e.g. emergent, floating and submerged), physical structure (e.g. shaded, open, shallow and deep water) and combinations thereof. For aquatic plants, 25 2x2 m quadrats were sampled in each wetland, distributed across these mesohabitats with all plants (i.e. all submerged, floating-leaved, emergent and marginal plants, including tree saplings and bryophytes) identified to the highest feasible taxonomic level. Nomenclature followed the Finnish Biodiversity Information Facility (FinBIF, <https://laji.fi/en>). Abundance per quadrat was estimated visually based on a 0-100% cover scale. For beetles, six 2x2 m quadrats were split among the major mesohabitats per wetland. Using a pond net (GB net with 1mm mesh) each quadrat was exhaustively sampled by vigorously sweeping through the vegetation, sediment and detritus until no obvious new species were found. Beetles were live-sorted and counted in the field with a few individuals retained as vouchers for species confirmation. Beetles were identified with the assistance of the following sources (Nilsson and Holmen 1995) and nomenclature followed (Vondel 2005, 2023; Przewoźny 2018; Nilsson and Hájek 2021, 2024).

eDNA Metabarcoding

Full details of the eDNA methods are provided in the Supporting Information. In brief, ten 1.5 L water samples were collected at roughly equidistant points around the

shoreline of each wetland. Samples were filtered through 0.45 µM PVDF Sterivex filter units (Millipore) until the filter clogged, and Longmire's solution (Longmire et al. 1997) then added as preservative. DNA was extracted using the Mu-DNA extraction protocol (Sellers et al. 2018). A filtration blank of distilled water and an extraction blank was included for each wetland. DNA was PCR-amplified in triplicate, in a two-step protocol, using two primer sets that have been designed to target a 106bp fragment of the mitochondrial 12S ribosomal RNA gene in vertebrates (F1/R1) (Riaz et al. 2011; Kelly et al. 2014), and a 141bp region of the COI in benthic invertebrates (fwhF2/EPTDr2n) (Leese et al. 2021) respectively. PCR negatives were included for each 24-sample sub-library. PCR products were pooled, purified and quantified, and the final vertebrate and invertebrate libraries were sequenced separately at 13 pM on an Illumina MiSeq using a 2 x 300 bp V3 kit (Illumina).

A custom 12S reference database for Finland was created based on records from FinBIF and local expert knowledge. For invertebrates, an updated version of the reference database developed by Harper et al. (2019a) (<https://doi.org/10.5281/zenodo.3993125>) was used, including a wider UK species list extracted from Freshbase (<https://freshbase.myspecies.info/>). Following the process outlined in Griffith et al. (2024), priority was given to species records present on the Barcode of Life Data System (BOLD) (<https://www.boldsystems.org/index.php/>), before checking any missing records for presence in GenBank. The final BLAST database was created from the reference sequences using makeblastdb (<https://www.ncbi.nlm.nih.gov/books/NBK569841/>) and included the accession-to-taxid map (-taxid_map) to allow for taxonomic assignment of OTUs for downstream Tapirs analysis (Griffith et al. 2024).

Bioinformatics steps, from quality control to taxonomic assignment, were performed using Tapirs, a reproducible workflow for the analysis of DNA metabarcoding data (<https://github.com/EvoHull/Tapirs>), with full details provided in the Supplementary Material. Following taxonomic assignment, taxa with <5 reads (vertebrates) or <10 reads (invertebrates) were removed to minimise noise based on inspection of read frequency distributions. A contaminant-specific threshold per sub-library was applied i.e. taxa were removed when the read count was below those found in site blanks. Reads from humans, domestic species, and species absent from the Evo region, were removed from the dataset (details in the Supporting Information). *Castor canadensis* reads were also removed so estimates of mammal species richness were not inflated. For analysis, four invertebrate groups (segmented worms (Lumbriculida and Tubificida), microcrustacea (Diplostraca and Cyclopoida), true flies (Diptera) and may/stone/caddisflies (Ephemeroptera, Plecoptera and Trichoptera)) which collectively represented 95% of invertebrate eDNA detections were retained, alongside four vertebrate species groups (amphibians, fish, birds and mammals). Vertebrates were assigned to species level where possible. For invertebrate eDNA samples, we used genus level resolution to minimize bias from incomplete reference database coverage at species level. Each invertebrate genus was checked on FinBIF to confirm their presence in southern central Finland.

Data treatment, exploration and analyses

Water physico-chemistry data (size, alkalinity, chlorophyll A, DOC, pH, total nitrogen and total phosphorus) were scaled and centred, with relationships among them observed using principal components analysis (PCA). Species richness was defined as the number of

species (or next highest taxonomic resolution individuals if reads could not be identified to species- or genus-level) or, for eDNA-derived invertebrate data, number of genera. Alpha diversity is defined as the number of taxa in that taxonomic group per site, and gamma diversity as the number of taxa in that group found in the landscape. Using the library lmerTest (Kuznetsova et al. 2016) taxa richness differences between wetland types at both sample and site scales were tested, initially, using generalised linear models with Poisson error distributions to account for count data. At the sample scale, each model was then tested for spatial dependence using Moran's I statistic. Spatial dependence was detected for plants (Moran's I = 0.232, $p = 0.001$), beetles (Moran's I = 0.086, $p = 0.044$), fish (Moran's I = 0.132, $p = 0.017$), amphibians (Moran's I = 0.170, $p = 0.008$). Spatial dependence was not detected for segmented worms (Moran's I = 0.060, $p = 0.126$), microcrustaceans (Moran's I = 0.035, $p = 0.200$), true flies (Moran's I = -0.05, $p = 0.829$), may/stone/caddisflies (Moran's I = 0.074, $p = 0.099$), birds (Moran's I = -0.022, $p = 0.607$) or mammals (Moran's I = 0.009, $p = 0.352$). Therefore, to account for the spatially nested experimental design of the study, we used generalised linear mixed models with a Poisson error distribution with site name as the random effect. All models were checked for overdispersion, but none was found.

Indicator taxa in each taxonomic group (based on occurrence and relative abundance or presence of taxa per wetland type) were derived via the 'multipatt' function from the R package 'indicspecies' (Cáceres and Legendre 2009). For beetles and eDNA samples, indicator taxa analysis was based on presence/absence matrices. Data on rarity and conservation status of each taxon (e.g. Status in Finland, Finnish Red List Category and Regulatory status) was extracted from the FinBIF.

To standardise taxa accumulation rates per wetland type, rarefaction curves were calculated for each of the 10 taxonomic groups based on incidence data per sample, except

for beetles, which were rarified based on the number of individuals per sweep. To assess differences in composition between wetland types (beta diversity), we used non-metric multidimensional scaling (NMDS) with a binomial dissimilarity index (presence only) for each taxon group. Except for aquatic plants, where a Bray-Curtis dissimilarity index was applied due to confidence in these abundance data. A permutational analysis of variance was used to test for differences in composition between wetland types within each taxon group using the function 'adonis2' (Oksanen et al. 2019). Furthermore, to check for spatial dependency, we compared the pairwise dissimilarity and Euclidean matrices for each species group using Mantel tests.

Ecological preference and functional diversity

We focussed on aquatic plants only as they respond most directly to beaver-induced habitat changes and are of fundamental importance to wider biodiversity, whether as a resource or component of habitat structure. In addition, limited trait data were available for the other taxon groups, or they showed minimal variation due to low taxa numbers e.g. amphibians, birds and mammals. To test for plant functional differences between each wetland type, we selected preferences for light, moisture, nitrogen (fertility or productivity) and reaction (soil acidity/pH) based on Ellenberg scores, obtained from Schmidt-Kloiber and Hering (2015). These preferences were chosen as they are well-described and understood, should reflect the variation in physico-chemical conditions experienced and the corresponding physiological or morphological traits of those species and are hence likely to indicate wider ecosystem function. Before functional diversity was calculated these quantitative preferences were rescaled to zero mean and standard deviation of one (z-

score). Using the R package FD (Laliberté, E. et al. 2014) we calculated functional richness, evenness and dispersion per site with plant abundance included. These represent the three dimensions of functional diversity. Firstly, richness - how much total diversity exists within a dataset). Secondly, evenness - bounded between 0 and 1, a measure of how regularly the species are situated along the trait space. Evenness is close to 0 when most species (and abundances) are tightly packed in a portion of the trait space, while it is close to 1 if species are regularly spread (with even abundances). Thirdly, dispersion (how abundance is spread among the different traits) (Villéger et al. 2008; Pavoine and Bonsall 2011). To explain which ecological preferences were driving functional differences between wetland types, we calculated Community-Weighted Means (CWM) for each trait also using the FD package.

All statistical analyses and graphics were generated using R Studio version 2024.04.1 (R Core Team 2022) with the additional packages; iNEXT (Hsieh et al. 2016), ggplot2 (Wickham and Chang 2016) and vegan (Oksanen et al. 2019).

Results

Both beaver and control wetlands had broadly similar physico-chemical characteristics (Table S2); mesotrophic, slightly acidic with high DOC, reflecting their location within a boreal forest landscape. From the principal components analysis (PCA), a large overlap among sites was apparent regardless of wetland type, with the first two axes explaining >90% of the variation, indicative of high similarity in physico-chemical conditions among sites (Fig. S1). Beaver wetlands were generally larger (2.1 ± 0.6 ha (0.7-4.1)) than control wetlands (1.2 ± 0.2 ha (0.5-2.5)), mean \pm SE (range), but not statistically so.

Alpha, beta and gamma diversity

Across the 18 wetlands surveyed, 380 taxa were detected using in situ (102 plant and 80 beetle species) and eDNA surveys (5 segmented worm, 13 microcrustacea, 114 true fly, 13 may/stone/caddisfly genera, and 4 amphibian, 10 fish, 25 bird and 14 mammal species), with 316 taxa overall being found in beaver wetlands and 274 in control wetlands (a full species inventory per site and taxonomic group is available in Supplementary Information Table S3).

Following quality control, taxonomic assignment, and threshold application and removal of non-target taxa, the final vertebrate eDNA metabarcoding dataset consisted of 7,216,204 reads over 175 samples (41,235 average read depth per sample). For invertebrate eDNA, the dataset consisted of 8,135,253 reads from 177 samples, with an average read depth of 45,962 per sample after non-target taxa had been removed. *C. canadensis* was detected in 26/90 samples from beaver wetlands at high read count (mean = 3906). By contrast, *C. canadensis* was only detected in 2/89 samples from control sites, at very low read count (mean = 190).

Per taxonomic group, the mean and total richness at sample and site level were mostly similar between wetland types (Table 1, Figs. S2 and S3). The only exceptions to this were plants and true flies, which had significantly higher richness in beaver wetlands at sample (true flies) and site-scale (plants and true flies). For all taxonomic groups, a large percentage of taxa were shared among wetland types (mean = 63%, range = 40-100%), but both beaver and control wetlands had unique taxa, mean = 19% (range = 0-40%) and mean = 18% (range = 0-40%), respectively (Fig. 3).

Indicator taxa were common for both plants and beetles in each wetland type (full list in Table S4). For plants, species from both habitat types were typical of northern latitude freshwater wetlands and peatlands, though species found in beaver wetlands were typical of either more disturbed conditions (*Alisma plantago-aquatica*, *Callitriche* spp.), unpalatable to beavers (*Cicuta virosa*, *Hippuris vulgaris*) or reflected the diversity of water depth and fluctuations (e.g. *Glyceria fluitans*, *Typha latifolia*). Plants in control wetlands were indicative of stable hydrological conditions (*Andromeda polifolia* and *Scheuchzeria palustris*) and saturated peat or deep water (*Vaccinium oxycoccos*, *Nuphar lutea*). Beetle species found across both wetland types were broadly indicative of mesotrophic, acidic sites with peat and *Sphagnum* substrates (e.g. *Ilybius aenescens*, *Ilybius guttiger*, *Bidessus grossepunctatus* and *Enochrus affinis*). Beetles associated with beaver wetlands were indicative of stagnant, small and well-vegetated wetlands (e.g. *Haliplus heydeni*, *H. ruficollis*, *Ilybius fuliginosus*, *I. quadriguttatus*, *Rhantus exsoletus*, *Hydroporus dorsalis* and *Hygrotus inaequalis*), whereas in control wetlands were primarily indicative of bryophyte-dominated bogs (*Hydroporus neglectus*, *H. scalesianus*, *H. tristis*, *Enochrus affinis* and *Contacyphon kongsbergensis*). Beetles were more commonly found in the shallow margins of the control wetlands rather than open water, whereas beetles were consistently found across the whole beaver wetland. Several true fly genera within the families of Chironomidae and Culicidae were indicators of beaver wetlands, but overall, we found fewer indicator taxa from eDNA samples. However, differences in occupancy (proportion of samples in which a taxon was detected) occurred (Figs. S4-S13), particularly for northern pike (*Esox lucius*) and ducks (Anatidae) both of which had consistently higher occupancy in beaver (84% and 78% of samples, respectively) compared to control wetlands (43% and 48%, respectively), but were not statistically significant.

Table 1. Summary of biodiversity metrics per taxonomic group per wetland type.

Taxonomic group	Wetland type (no. of samples)	Mean sample taxa \pm SE (min- max)	Mean site taxa \pm SE (min- max)	Total taxa observed	Indicator taxa
Plants	Beaver (n = 225)	7.6 \pm 0.2 (1 – 15)	38.4 \pm 1.2 (35 – 45)	92	37
	Control (n = 225)	7.3 \pm 0.2 (1 – 13)	26.0 \pm 1.9 (17 – 34)	68	20
Beetles	Beaver (n = 54)	5.9 \pm 0.4 (1 – 15)	20.2 \pm 2.0 (12 – 30)	58	8
	Control (n = 54)	6.2 \pm 0.4 (1 – 15)	19.8 \pm 1.5 (14 – 26)	65	6
Segmented worms	Beaver (n = 90)	1.7 \pm 0.1 (1 – 4)	3.1 \pm 0.5 (2 – 5)	5	0
	Control (n = 90)	1.4 \pm 0.1 (1 – 4)	2.3 \pm 0.5 (1 – 4)	5	0
Microcrustacea	Beaver (n = 90)	2.5 \pm 0.1 (1 – 6)	5.9 \pm 0.6 (3 – 8)	11	1
	Control (n = 90)	2.7 \pm 0.2 (1 – 6)	5.6 \pm 0.9 (2 – 10)	13	1
True flies	Beaver (n = 90)	14.3 \pm 0.8 (3 – 31)	38.6 \pm 4.3 (24 – 58)	100	19
	Control (n = 90)	9.8 \pm 0.5 (1 – 23)	27.2 \pm 2.8 (17 – 43)	75	2
May/Stone/Caddisflies	Beaver (n = 90)	1.9 \pm 0.2 (1 – 7)	4.6 \pm 0.9 (1 – 10)	12	1

	Control (n = 90)	1.7 ± 0.2 (1 – 4)	2.7 ± 0.8 (1 – 6)	8	0
Amphibians	Beaver (n = 90)	1.2 ± 0.1 (1 – 2)	2.0 ± 0.4 (1 – 3)	4	0
	Control (n = 90)	1.2 ± 0.1 (1 – 3)	1.8 ± 0.3 (1 – 3)	4	0
Fish	Beaver (n = 90)	2.2 ± 0.1 (1 – 5)	4.0 ± 0.5 (2 – 7)	7	0
	Control (n = 90)	2.4 ± 0.1 (1 – 4)	4.1 ± 0.4 (3 – 6)	9	1
Birds	Beaver (n = 90)	1.5 ± 0.1 (1 – 5)	4.0 ± 0.8 (1 – 8)	17	0
	Control (n = 90)	1.7 ± 0.2 (1 – 8)	3.9 ± 0.8 (2 – 10)	18	0
Mammals	Beaver (n = 90)	1.4 ± 0.2 (1 – 5)	2.4 ± 0.5 (1 – 5)	10	0
	Control (n = 90)	1.3 ± 0.1 (1 – 2)	2.7 ± 0.8 (1 – 5)	9	0




Figure 3. Ven diagrams of taxa richness and percentage of taxa unique to each wetland type; beaver (blue, left circles) and control (orange, right circles) per taxonomic group. Areas of overlap represent shared taxa.

No nationally rare plant species were found in the study. *Elodea canadensis* was the only non-native plant species recorded, though this only occurred in one quadrat. The beetles *Dytiscus latissimus* and *Graphoderus bilineatus* were found in two and one beaver wetland, respectively, with individuals identified visually and returned. Both are named on the Bern Convention and listed as vulnerable on the EU habitats directive (Foster 1996b, a). Genera within the segmented worms, microcystacea, true flies and may/stone/caddisflies were generally common and widespread. All amphibian and fish species found were classed as 'Least Concern' under IUCN Red List status, however all amphibians are protected within Finland under national (Nature Conservation Act) or EU law (EU Habitats Directive). No local or internationally rare birds or mammals were found.

Accumulation curves (Fig. 4) implied that sampling coverage was high (mean = 96%) for most taxonomic groups, though curves for true flies, birds and mammals indicated that increasing the sampling effort would substantially increase the number of taxa recorded (Fig. 4 E, I and J). Overall, rates of taxa accumulation were similar between wetland types, with only plants and true flies accumulating at a noticeably greater rate in beaver and control wetlands, respectively.




Figure 4. Accumulation curves for beaver (blue) and control wetlands (orange lines) per taxonomic group; a) aquatic plants, b) beetles, c) segmented worms, d) microcrustaceans, e) true flies, f) may/stone/caddisflies, g) amphibians, h) fish, i) birds and j) mammals. Lines represent sample-based accumulation at sample scale for all taxonomic groups, apart from beetles which were individual-based.

Overlaps were observed in all taxonomic groups, indicating shared taxa between wetland types. However, there were clear and significant differences in the assemblages of most taxonomic groups between wetland types (Fig. 5). No significant differences between wetland types were found for segmented worms, amphibians or mammals (Fig. 5 C, G and J). Spatial autocorrelation via Mantel tests was not detected for any of the taxonomic groups; plants ($r = -0.02$, $p = 1$), beetles ($r = -0.02$, $p = 0.929$), segmented worms ($r = -0.008$, $p = 0.736$), microcrustaceans ($r = -0.032$, $p = 1$), true flies ($r = -0.024$, $p = 1$), may/stone/caddisflies ($r = -0.024$, $p = 0.919$), amphibians ($r = -0.007$, $p = 0.620$), fish ($r = -0.034$, $p = 1$), birds ($r = -0.001$, $p = 0.486$) or mammals ($r = 0.006$, $p = 0.450$).




Figure 5. Non-metric multidimensional scaling (NMS) unconstrained ordination plots for a) aquatic plants, b) beetles, c) segmented worms, d) microcrustaceans, e) true flies, f) may/stone/caddisflies, g) amphibians, h) fish, i) birds and j) mammals. Each panel contains a spider plot for; beaver (blue) and control (orange) wetlands, with each sample connected to the waterbody type centroid. P-values represent outputs from a permutational multivariate analysis of variance.

Functional diversity

Plants in beaver wetlands had greater functional richness, evenness and dispersion compared to control wetlands (Fig. 6 a, b, c), significantly so for richness. This indicates that a greater combination of functions (i.e. ecological preferences) was present in beaver wetlands reflecting the greater variation in habitat conditions among patches within these wetlands compared to control wetlands. Of the ecological preferences used to obtain functional diversity scores, beaver wetlands also had lower CWM scores for light, higher for moisture and significantly higher for nitrogen (Fig. 6 d, e and f). This signifies that vegetation found in beaver wetlands was indicative of tolerance for more shaded, moister and fertile conditions.




Figure 6. Error plots comparing aquatic plant functional a) richness, b) evenness, c) dispersion, and Community-Weighted Means values for d) light, e) moisture and f) nitrogen for beaver (blue) and control wetlands (orange). Coloured points represent the mean of the data, error bars are equal to one standard error and grey dots show individual data points. P values are outputs from Benjamini-Hochberg corrected general linear models that tested differences between wetland types per function or trait.

Discussion

Ecosystem engineers are recognised to stimulate environmental heterogeneity and biodiversity, warranting their role in ecological restoration. With regards to beavers, recent studies highlight that these benefits can also spillover ‘beyond the pond’ (Orazi et al. 2022;

Fedyń et al. 2023; Hooker et al. 2024; Cook et al. 2025). However, very few studies have demonstrated that these patterns apply simultaneously across multiple taxa groups and sites in parallel, and within a common landscape. We found that beaver-created wetlands were both speciose and compositionally distinct at the landscape-scale for some taxonomic groups and added to the regional taxon pool for most of these groups. We also found that major biotic differences between beaver-created and control wetlands at the bottom of the food chain (i.e. aquatic plants, microcrustacea and benthic invertebrates) imply the potential to reconfigure trophic networks.

What do beaver wetlands add to a landscape?

Despite finding minor differences in physico-chemistry between wetland types, major physical differences were apparent between wetland types. Control wetlands were hydrologically stable with steep margins typical of glacial kettle hole lakes. They were broadly circular and deep in the middle, moving from a saturated marginal zone to deep water (>1 m) over a short distance that restricts emergent vegetation (Nummi et al. 2019a). Beaver wetlands had relatively shallow, complex margins with highly variable depths (Hood and Larson 2014) (the deepest section most often being the original stream or drainage channel). Physical variation was further exemplified by the 'mosaic of mess' beavers create e.g. selectively felled trees, remnants of grazed or uprooted shrubs and aquatic plants, with woody debris – coarse, fine and (standing or fallen) dead wood – scattered across the wetland (Thompson et al. 2016). Furthermore, there was evidence of beavers excavating canals and maintaining their dams by moving sediment, stones and wood (Larsen et al. 2021). Individually, these dynamic processes are difficult to quantify, or replicate at suitable

scales via conventional interventionist approaches, but collectively they enhance within-wetland heterogeneity that drives biodiversity (Willby et al. 2018).

Though habitat patches were often visually different between wetland types, the biodiversity benefits of beaver wetlands were not always obvious at small scales. For example, richness for each taxonomic group was similar between wetland types, particularly at sample level. Biodiversity differences accrued through the cumulative collection of multiple samples that were highly heterogenous, particularly for aquatic plants and true flies. We found significant compositional differences between wetland types for aquatic plants, beetles, microcrustacea, true flies, may/stone/caddisflies, fish and birds, but not segmented worms, amphibians and mammals. This most likely reflected the small species pool of amphibians in the study area and small number of worms and mammals detected via eDNA. Across the taxonomic groups, on average 61% of taxa were found in both wetland types, but each type contained unique and indicative species. For example, in beaver wetlands, plants were indicative of fluctuating water levels and higher fertility, while beetles were indicative of smaller sites and pools (Foster et al. 2016, 2020), which likely reflects the greater habitat complexity and heterogeneity in depth in beaver wetlands, creating predator and prey refugia. Due to the high within-site heterogeneity in beaver wetlands and physical differences from control wetlands, the addition of beaver wetlands to the landscape increased freshwater biodiversity by an average of 19% (range 0-40%) across all groups. In Evo, Finland the landscape is already rich in relatively undisturbed waterbodies, yet we found biodiversity gains similar to beaver wetlands in Sweden (~30% for aquatic plants and beetles) that were surrounded by agricultural, forestry or suburban landuses (Law et al. 2019).

Sampling biodiversity of wetlands using eDNA is complementary to in-situ surveys and allows detection of more mobile or elusive taxa and is indicative of recent species: water interactions (Harper et al. 2019b). A key finding from this sampling approach was the greater occupancy of ducks and pike in beaver compared to control wetlands. Previous research in this geographical region has reported the favourability of beaver wetlands for teal and other dabbling duck species due to food and habitat provisioning (Nummi and Hahtola 2008; Nummi and Holopainen 2014). More broadly, a greater abundance and differing composition of birds was reported from around Polish beaver wetlands (Fedyń et al. 2023). Many net sweeps we used to sample beetles also captured juvenile pike <10 cm (A. Law, *pers. obs.*) suggesting that this age class benefits from the shallow and well-vegetated wetland margins that are a feature of beaver wetlands but absent from other freshwater habitats in this landscape (Nummi and Hahtola 2008).

As the rarefaction curves did not reach an asymptote for most taxonomic groups (excluding worms and amphibians), further taxa are present in the landscape that were not detected by our sampling methods. For eDNA, our findings are additionally influenced by taxa ecology and survey timing. Further species would be detected with seasonally replicated campaigns, particularly for waterfowl that nest early in the year and disperse by late-summer (Nummi & Pöysä, 1993). Mammal detection (semi-aquatic, arboreal and terrestrial) using eDNA is highly variable across surveyed sites due to spatial and temporal clustering (Harper et al. 2019b) i.e. reduced species: water interactions as there are fewer, larger but mobile taxa in the landscape. However, it offers a useful complementary survey approach to camera trapping and footprint surveys which have previously served to demonstrate the unique value of beaver wetlands in Evo to mammals (Nummi et al. 2019a). Given that beaver wetlands are highly heterogeneous, taxa occurrences detected via eDNA

may be even more localised, particularly in comparison to control wetlands that have a more homogeneous physical environment. Overall, similar trends were evident for each sampling technique, i.e. a distinct taxonomic composition between wetland types (except worms, amphibians and mammals), which could have knock-on effects for both aquatic and terrestrial trophic networks (Dewey et al. 2025).

Taxonomic and functional metrics are complementary (Lu et al. 2023) but can elucidate different mechanisms and insights. Despite this, studies comparing functional diversity between wetland types are rare (though growth strategies were explored in (Law et al. 2019)). Our finding that beaver wetlands had a higher functional richness demonstrates that aquatic plants in beaver wetlands are indicative of multiple ecological preferences with minimal redundancy. The driver of increased function is likely related to niche differentiation driven by high local habitat heterogeneity. Since succession and disturbance also affect functional diversity (Coccia et al. 2024) we recognise that some of the observed increase in functional diversity will be driven by ruderal species that colonise quickly, have broad environmental tolerances and are weak competitors (Ruhí et al. 2013), their habitats being renewed by disturbance events arising from beaver activity. Overall, our results demonstrate that beaver wetlands not only improve landscape biodiversity they also increase landscape ecosystem resilience through increased functional diversity.

Implications and research gaps

Several studies on beaver wetland biodiversity have indicated increased richness at alpha and gamma scales (see (Stringer and Gaywood 2016) for a review). However, for this study and many other restoration and rewilding studies, a key question remains: where do

these additional taxa come from? We propose 3 explanations for beaver-modified landscapes. Firstly, a spatially and temporally hierarchical disturbance mosaic, e.g. at fine scales, tree felling and selective herbaceous grazing, at coarse scales water level fluctuations due to dam construction and degradation, or decadal scale cycles of abandonment and reoccupation. This creates dynamic environments which limits highly competitive taxa and creates unique opportunities for weakly-competitive taxa to persist indefinitely within the landscape (Kivinen et al. 2020). Secondly, flooding and reflooding, combined with local scale digging and uprooting, will expose and awaken dormant seedbanks (Abernethy and Willby 1999; Law et al. 2017; Alderton et al. 2017), some potentially in long-abandoned beaver wetlands which may be re-occupied. Thirdly, increased abundance and quality of freshwater in the landscape increases and diversifies connectivity pathways leading to more stochastic colonisation events (Baastrup-Spohr et al. 2016), as has been demonstrated within the Evo landscape (Kivinen and Nummi 2025). For example, due to habitat productivity (invertebrate biomass) and structure (large areas of shallow water) waterbird richness is higher in beaver-created wetlands than other types (Nummi and Hahtola 2008; Nummi and Holopainen 2014). In promoting connectivity, beaver wetlands should therefore facilitate seed dispersal by waterfowl (Baastrup-Spohr et al. 2016).

Conclusions

The restoration of modern landscapes via rewilding has captivated conservationists and the public globally as a nature-based, sustainable and passive approach (Pettorelli et al. 2018). Given the biodiversity crisis and pressing targets to arrest biodiversity loss, the complete toolbox of restoration methods must be utilised to kickstart ecosystem processes

that will bring benefits at scale. We provide empirical evidence that a major ecosystem engineer can improve biodiversity both within and beyond its wetland environment across multiple taxonomic groups within the same uniform landscape. Our results demonstrate that beaver-created wetlands will support taxa absent from the closest comparable habitats in this landscape and are integral to improving landscape functionality. Following near extirpation in the early 1900s after centuries of hunting the geographical range of beavers is expanding across North America and Europe, aided by legal protection, reintroductions and translocations having reached about 10% of its historical maximum. Viewed in this context, our findings hint at the likely scale of past biodiversity loss associated with beaver-dependent wetlands, while offering a glimpse of what could be gained from their ongoing population recovery.

Acknowledgements

We thank staff at Lammi Biological Station (University of Helsinki) for their logistical and technical support, in addition to interns who helped with water filtration, Charlotte Buseti, Aleya Nur Erim and Sophie Louise Stafford. We also thank Bernd Haenfling, Lucio Marcello and Nathan Griffiths (University of Highlands and Islands) for provision of invertebrate eDNA databases. Comments by two anonymous reviewers helped improve this manuscript.

Statements and declarations

Funding. Funding from Carnegie Trust for the Universities of Scotland enabled AL, NW, TS and DB to carry out fieldwork in Finland. Funding from Biotechnology and Biological Sciences Research Council (BBSRC) allowed a follow-up workshop in Finland for all authors

to complete the first manuscript draft. Natural England co-funded TS's PhD, and Hydronations for DB's PhD.

Competing interests. The authors have no relevant financial or non-financial interests to disclose.

Ethical approval was sought and obtained for the University of Stirling Animal Welfare and Ethical Review Body prior to data collection.

Author contributions.

All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by all authors. The first draft of the manuscript was written by Alan Law and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Data availability statement. Site-level data supporting the findings of this study are available within the paper and its Supplementary Information. Sample-level data that support the findings of this study are not openly available due to reasons of sensitivity and are available from the corresponding author upon reasonable request.

References

Abernethy VJ, Willby NJ (1999) Changes along a disturbance gradient in the density and composition of propagule banks in floodplain aquatic habitats. *Plant Ecology* 140:177–190

Alberdi A, Aizpurua O, Gilbert MTP, Bohmann K (2018) Scrutinizing key steps for reliable metabarcoding of environmental samples. *Methods in Ecology and Evolution* 9:134–147. <https://doi.org/10.1111/2041-210X.12849>

Alderton E, Sayer CD, Davies R, et al (2017) Buried alive: Aquatic plants survive in 'ghost ponds' under agricultural fields. *Biological Conservation* 212:105–110. <https://doi.org/10.1016/j.biocon.2017.06.004>

- Andersen LH, Nummi P, Bahrndorff S (2024) Can beavers help improve terrestrial invertebrate diversity? *Front Ecol Evol* 12:.
<https://doi.org/10.3389/fevo.2024.1396207>
- Andersen LH, Ransborg C, Pertoldi C, et al (2023) Can reintroduction of beavers improve insect biodiversity? *Journal of Environmental Management* 337:117719.
<https://doi.org/10.1016/j.jenvman.2023.117719>
- Araújo MB, Alagador D (2024) Expanding European protected areas through rewilding. *Current Biology* 0: <https://doi.org/10.1016/j.cub.2024.07.045>
- Arvola L, Rask M, Huotari J, et al (2025) Chemical responses of small boreal lakes to atmospheric and catchment drivers over four decades. *Science of The Total Environment* 968:178696. <https://doi.org/10.1016/j.scitotenv.2025.178696>
- Baastrup-Spohr L, Kragh T, Petersen K, et al (2016) Remarkable richness of aquatic macrophytes in 3-years old re-established Lake Fil, Denmark. *Ecological Engineering* 95:375–383. <https://doi.org/10.1016/j.ecoleng.2016.06.081>
- Brazier RE, Puttock A, Graham HA, et al (2020) Beaver: Nature’s ecosystem engineers. *WIREs WATER* e1494:. <https://doi.org/10.1002/wat2.1494>
- Burchsted D, Daniels M, Thorson R, Vokoun J (2010) The river discontinuum: applying beaver modifications to baseline conditions for restoration of forested headwaters. *BioScience* 60:908–922. <https://doi.org/10.1525/bio.2010.60.11.7>
- Bush BM, Wissinger SA (2016) Invertebrates in beaver-created wetlands and ponds. In: *Invertebrates in Freshwater Wetlands: An International Perspective on Their Ecology*. pp 411–449
- Butler DR, Malanson GP (1995) Sedimentation rates and patterns in beaver ponds in a mountain environment. *Geomorphology* 13:255–269
- Byers JE (2022) Using ecosystem engineers to enhance multiple ecosystem processes. *Functional Ecology* 38:22–36. <https://doi.org/10.1111/1365-2435.14130>
- Byers JE, Cuddington K, Jones CG, et al (2006) Using ecosystem engineers to restore ecological systems. *Trends in Ecology & Evolution* 21:493–500.
<https://doi.org/10.1016/j.tree.2006.06.002>
- Cáceres MD, Legendre P (2009) Associations between species and groups of sites: indices and statistical inference. *Ecology* 90:3566–3574. <https://doi.org/10.1890/08-1823.1>
- Chen S, Zhou Y, Chen Y, Gu J (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* 34:i884–i890. <https://doi.org/10.1093/bioinformatics/bty560>
- Ciechanowski M, Kubic W, Rynkiewicz A, Zwolicki A (2011) Reintroduction of beaver *Castor fiber* may improve habitat quality for vespertilionid bats foraging in small river valleys. *European Journal of Wildlife Research* 57:737–747.
<https://doi.org/10.1007/s10344-010-0481-y>

- Coccia C, Almeida BA, Badosa A, et al (2024) Hydroperiod length, not pond age, determines zooplankton taxonomic and functional diversity in temporary ponds. *Ecological Indicators* 159:111632. <https://doi.org/10.1016/j.ecolind.2024.111632>
- Coccia C, Almeida BA, Green AJ, et al (2021) Functional diversity of macroinvertebrates as a tool to evaluate wetland restoration. *Journal of Applied Ecology* 58:2999–3011. <https://doi.org/10.1111/1365-2664.14038>
- Cook P, Law A, Pattison Z, Willby NJ (2025) Beaver wetlands create a buzz and a flutter for pollinators. *Journal of Applied Ecology* 62:3288–3299. <https://doi.org/10.1111/1365-2664.70193>
- Dewey L, Fritz KA, Kirschman LJ, et al (2025) Beaver recolonization explains aquatic insect emergence patterns. *Freshwater Science* 000–000. <https://doi.org/10.1086/734565>
- Dudgeon D (2019) Multiple threats imperil freshwater biodiversity in the Anthropocene. *Current Biology* 29:R960–R967. <https://doi.org/10.1016/j.cub.2019.08.002>
- Fedyń I, Przepióra F, Sobociński W, et al (2023) Beyond beaver wetlands: The engineering activities of a semi-aquatic mammal mediate the species richness and abundance of terrestrial birds wintering in a temperate forest. *Forest Ecology and Management* 529:120698. <https://doi.org/10.1016/j.foreco.2022.120698>
- Fedyń I, Sobociński W, Czyżowicz S, et al (2024) Ecosystem engineers cause biodiversity spill-over: Beavers are associated with breeding bird assemblages on both wetlands and adjacent terrestrial habitats. *Science of The Total Environment* 950:175166. <https://doi.org/10.1016/j.scitotenv.2024.175166>
- Foster G (1996a) *Dytiscus latissimus*. The IUCN Red List of Threatened Species. IUCN Red List of Threatened Species
- Foster G (1996b) *Graphoderus bilineatus*. The IUCN Red List of Threatened Species. IUCN Red List of Threatened Species
- Foster GN, Bilton DT, Hammond M, Nelson BH (2020) Atlas of water beetles of Britain and Ireland - smaller families of polyphaga. Biological Records Centre
- Foster GN, Bilton DT, Nelson BH (2016) Atlas of the predaceous water beetles (Hydradephaga) of Britain and Ireland. Field Studies Council
- Gardner RC, Finlayson C (2018) Global Wetland Outlook: State of the World's Wetlands and Their Services to People. Social Science Research Network, Rochester, NY
- Gaywood MJ, Stanley-Price M (2022) Moving Species: Reintroductions and Other Conservation Translocations. In: Moehrenschrager A, Ewen JG, Gaywood MJ, Hollingsworth PM (eds) Conservation Translocations. Cambridge University Press, Cambridge, pp 3–42
- Griffith N, Hänfling B, Andreou D, et al (2024) eDNA monitoring of restoration effects on the River Severn (Phase II) Unlocking the Severn LIFE project. Natural England

- Harper LR, Griffiths NP, Lawson Handley L, et al (2019a) Development and application of environmental DNA surveillance for the threatened crucian carp (*Carassius carassius*). *Freshwater Biology* 64:93–107. <https://doi.org/10.1111/fwb.13197>
- Harper LR, Lawson Handley L, Carpenter AI, et al (2019b) Environmental DNA (eDNA) metabarcoding of pond water as a tool to survey conservation and management priority mammals. *Biological Conservation* 238:546218. <https://doi.org/10.1016/j.biocon.2019.108225>
- Hood GA, Larson DG (2013) Beaver-created habitat heterogeneity influences aquatic invertebrate assemblages in boreal Canada. *Wetlands* 34:19–29. <https://doi.org/10.1007/s13157-013-0476-z>
- Hood GA, Larson DG (2014) Ecological engineering and aquatic connectivity: a new perspective from beaver-modified wetlands. *Freshwater Biology* 60:198–208. <https://doi.org/10.1111/fwb.12487>
- Hooker J, Foxley T, Stone EmmaL, Lintott PaulR (2024) Re-establishing historic ecosystem links through targeted species reintroduction: Beaver-mediated wetlands support increased bat activity. *Science of The Total Environment* 951:175661. <https://doi.org/10.1016/j.scitotenv.2024.175661>
- Hsieh TC, Ma KH, Chao A (2016) iNEXT: An R package for rarefaction and extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution* 7:1451–1456. <https://doi.org/10.1111/2041-210X.12613>
- Hyvönen T, Nummi P (2008) Habitat dynamics of beaver *Castor canadensis* at two spatial scales. *Wildlife Biology* 14:302–308. [https://doi.org/10.2981/0909-6396\(2008\)14%255B302:HDOBCC%255D2.0.CO;2](https://doi.org/10.2981/0909-6396(2008)14%255B302:HDOBCC%255D2.0.CO;2)
- Kelly RP, Port JA, Yamahara KM, et al (2014) Harnessing DNA to improve environmental management. *Science* 344:1455–1456. <https://doi.org/10.1126/science.1251156>
- Kitson JJN, Hahn C, Sands RJ, et al (2019) Detecting host–parasitoid interactions in an invasive Lepidopteran using nested tagging DNA metabarcoding. *Molecular Ecology* 28:471–483. <https://doi.org/10.1111/mec.14518>
- Kivinen S, Nummi P (2025) Immediate facilitation and engineering legacy of beavers: 54 years of patch dynamics in a boreal landscape. *Science of The Total Environment* 999:180341. <https://doi.org/10.1016/j.scitotenv.2025.180341>
- Kivinen S, Nummi P, Kumpula T (2020) Beaver-induced spatiotemporal patch dynamics affect landscape-level environmental heterogeneity. *Environmental Research Letters* 15:. <https://doi.org/10.1088/1748-9326/ab9924>
- Köster J, Rahmann S (2012) Snakemake—a scalable bioinformatics workflow engine. *Bioinformatics* 28:2520–2522. <https://doi.org/10.1093/bioinformatics/bts480>

- Kuznetsova A, Brockhoff PB, Christensen RHB (2016) lmerTest: Tests in Linear Mixed Effects Models
- Laliberté, E., Legendre, P., Shipley, B. (2014) FD: measuring functional diversity from multiple traits, and other tools for functional ecology
- Larsen A, Larsen JR, Lane SN (2021) Dam builders and their works: Beaver influences on the structure and function of river corridor hydrology, geomorphology, biogeochemistry and ecosystems. *Earth-Science Reviews* 218:103623. <https://doi.org/10.1016/j.earscirev.2021.103623>
- Law A, Gaywood MJ, Jones KC, et al (2017) Using ecosystem engineers as tools in habitat restoration and rewilding: beaver and wetlands. *Science of the Total Environment* 605–606. <https://doi.org/10.1016/j.scitotenv.2017.06.173>
- Law A, Levanoni O, Foster G, et al (2019) Are beavers a solution to the freshwater biodiversity crisis? *Diversity and Distributions* 25:1763–1772. <https://doi.org/10.1111/ddi.12978>
- Law A, McLean F, Willby NJ (2016) Habitat engineering by beaver benefits aquatic biodiversity and ecosystem processes in agricultural streams. *Freshwater Biology* 61:486–499
- Leese F, Sander M, Buchner D, et al (2021) Improved freshwater macroinvertebrate detection from environmental DNA through minimized nontarget amplification. *Environmental DNA* 3:261–276. <https://doi.org/10.1002/edn3.177>
- Leung B, Hargreaves AL, Greenberg DA, et al (2020) Clustered versus catastrophic global vertebrate declines. *Nature* 588:267–271. <https://doi.org/10.1038/s41586-020-2920-6>
- Longmire JL, Maltbie M, Baker RJ (1997) Use of “Lysis Buffer” in DNA isolation and its implication for museum collections. 163:
- Lu K, Jähnig SC, Wu H, et al (2023) Trait-based approach of aquatic insects to track recovery of wetland ecosystems in Northeast China. *Ecological Indicators* 155:111012. <https://doi.org/10.1016/j.ecolind.2023.111012>
- Maasri A, Jähnig SC, Adamescu MC, et al (2022) A global agenda for advancing freshwater biodiversity research. *Ecology Letters* 25:255–263. <https://doi.org/10.1111/ele.13931>
- McCaffery M, Eby L (2016) Beaver activity increases aquatic subsidies to terrestrial consumers. *Freshwater Biology* 61:518–532. <https://doi.org/10.1111/fwb.12725>
- Moser V, Capitani L, Zehnder L, et al (2025) Habitat heterogeneity and food availability in beaver-engineered streams foster bat richness, activity and feeding. *Journal of Animal Ecology* 94:2403–2420. <https://doi.org/10.1111/1365-2656.70136>

- Mutillod C, Buisson É, Mahy G, et al (2024) Ecological restoration and rewilding: two approaches with complementary goals? *Biological Reviews* 99. <https://doi.org/10.1111/brv.13046>
- Nilsson AN, Hájek J (2021) A World Catalogue of the Family Dytiscidae, or the Diving Beetles (Coleoptera, Adephegata)
- Nilsson AN, Hájek J (2024) Catalogue of Palearctic Noteridae (Coleoptera)
- Nilsson AN, Holmen M (1995) The Aquatic Adephegata (Coleoptera) of Fennoscandia and Denmark, Volume II. Dytiscidae. E.J.Brill, Leiden
- Nummi P, Arzel C, Sauramo V (2021a) Populations in stable and variable habitats: Green and common sandpiper in a beaver-influenced landscape. *Global Ecology and Conservation* 28:e01678. <https://doi.org/10.1016/j.gecco.2021.e01678>
- Nummi P, Hahtola A (2008) The beaver as an ecosystem engineer facilitates teal breeding. *Ecography* 31:519–524. <https://doi.org/10.1111/j.0906-7590.2008.05477.x>
- Nummi P, Holopainen S (2020) Restoring wetland biodiversity using research: Whole-community facilitation by beaver as framework. *Aquatic Conservation: Marine and Freshwater Ecosystems* 30:1798–1802. <https://doi.org/10.1002/aqc.3341>
- Nummi P, Holopainen S (2014) Whole-community facilitation by beaver: ecosystem engineer increases waterbird diversity. *Aquatic Conservation: Marine and Freshwater Ecosystems* 24:326–633. <https://doi.org/10.1002/aqc.2437>
- Nummi P, Kattainen S, Ulander P, Hahtola A (2011) Bats benefit from beavers: a facilitative link between aquatic and terrestrial food webs. *Biodiversity and Conservation* 20:851–859. <https://doi.org/10.1007/s10531-010-9986-7>
- Nummi P, Kuuluvainen T (2013) Forest disturbance by an ecosystem engineer: beaver in boreal forest landscapes. *Boreal Environment Research* 18:13–24
- Nummi P, Liao W, Huet O, et al (2019a) The beaver facilitates species richness and abundance of terrestrial and semi-aquatic mammals. *Global Ecology and Conservation* e00701. <https://doi.org/10.1016/j.gecco.2019.e00701>
- Nummi P, Liao W, van der Schoor J, Loehr J (2021b) Beaver creates early successional hotspots for water beetles. *Biodivers Conserv* 30:2655–2670. <https://doi.org/10.1007/s10531-021-02213-8>
- Nummi P, Pöysä H (1993) Habitat associations of ducks during different phases of the breeding season. *Ecography* 16:319–328. <https://doi.org/10.1111/j.1600-0587.1993.tb00221.x>
- Nummi P, Suontakanen EM, Holopainen S, Väänänen VM (2019b) The effect of beaver facilitation on Common Teal: pairs and broods respond differently at the patch and landscape scales. *Ibis* 161:301–309. <https://doi.org/10.1111/ibi.12626>

- Oksanen AJ, Blanchet FG, Kindt R, et al (2019) *vegan: Community Ecology Package*
- Orazi V, Hagge J, Gossner MM, et al (2022) A Biodiversity Boost From the Eurasian Beaver (*Castor fiber*) in Germany's Oldest National Park. *Frontiers in Ecology and Evolution* 10:
- Palmer MA, Hondula KL, Koch BJ (2014) Ecological Restoration of Streams and Rivers: Shifting Strategies and Shifting Goals. *Annual Review of Ecology, Evolution, and Systematics* 45:247–269. <https://doi.org/10.1146/annurev-ecolsys-120213-091935>
- Parker H, Nummi P, Hartman G, Rosell F (2012) Invasive North American beaver *Castor canadensis* in Eurasia: a review of potential consequences and a strategy for eradication. *Wildlife Biology* 18:
- Pavoine S, Bonsall MB (2011) Measuring biodiversity to explain community assembly: a unified approach. *Biological Reviews* 86:792–812. <https://doi.org/10.1111/j.1469-185X.2010.00171.x>
- Pettorelli N, Barlow J, Stephens PA, et al (2018) Making rewilding fit for policy. *Journal of Applied Ecology* 55:1114–1125. <https://doi.org/10.1111/1365-2664.13082>
- Przewoźny M (2018) *Catalogue of Palearctic Hydrophiloidea (Coleoptera)*
- Puttock A, Graham HA, Ashe J, et al (2020) Beaver dams attenuate flow: A multi-site study. *Hydrological Processes* 1–18. <https://doi.org/10.1002/hyp.14017>
- Puttock A, Graham HA, Carless D, Brazier RE (2018) Sediment and nutrient storage in a beaver engineered wetland. *Earth Surface Processes and Landforms* 43:2358–2370. <https://doi.org/10.1002/esp.4398>
- R Core Team (2022) *R: A Language and Environment for Statistical Computing*
- Ray AM, Rebertus AJ, Ray HL (2001) Macrophyte succession in Minnesota beaver ponds. *Canadian Journal of Botany* 79:487–499
- Riaz T, Shehzad W, Viari A, et al (2011) *ecoPrimers: inference of new DNA barcode markers from whole genome sequence analysis*. *Nucleic Acids Research* 39:e145. <https://doi.org/10.1093/nar/gkr732>
- Rognes T, Flouri T, Nichols B, et al (2016) VSEARCH: a versatile open source tool for metagenomics. *PeerJ* 4:e2584. <https://doi.org/10.7717/peerj.2584>
- Rosell F, Campbell-Palmer R (2022) *Beavers: ecology, behaviour, conservation, and management*. Oxford University Press
- Ruhí A, Boix D, Gascón S, et al (2013) Nestedness and successional trajectories of macroinvertebrate assemblages in man-made wetlands. *Oecologia* 171:545–556. <https://doi.org/10.1007/s00442-012-2440-7>

- Schloemer S, Hörren T, Lorenz AW, Hering D (2023) The macroinvertebrate fauna of maintained and abandoned beaver dams. *Hydrobiologia* 850:1763–1778. <https://doi.org/10.1007/s10750-023-05176-9>
- Schmidt-Kloiber A, Hering D (2015) www.freshwaterecology.info – An online tool that unifies, standardises and codifies more than 20,000 European freshwater organisms and their ecological preferences. *Ecological Indicators* 53:271–282. <https://doi.org/10.1016/j.ecolind.2015.02.007>
- Sellers GS, Muri CD, Gómez A, Hänfling B (2018) Mu-DNA: a modular universal DNA extraction method adaptable for a wide range of sample types. *Metabarcoding and Metagenomics* 2:e24556. <https://doi.org/10.3897/mbmg.2.24556>
- Sinclair JS, Mademann JA, Haubrock PJ, Haase P (2023) Primarily neutral effects of river restoration on macroinvertebrates, macrophytes, and fishes after a decade of monitoring. *Restoration Ecology* 31:e13840. <https://doi.org/10.1111/rec.13840>
- Spyra A, Cieplok A, Krodkiewska M (2023) Beavers ecosystem altering: Influence of beaver dams on aquatic invertebrates in newly created beavers ponds and small mountain river. *Ecohydrology & Hydrobiology*. <https://doi.org/10.1016/j.ecohyd.2023.06.009>
- Stringer AP, Gaywood MJ (2016) The impacts of beavers *Castor* spp. on biodiversity and the ecological basis for their reintroduction to Scotland, UK. *Mammal Review* 46:270–283. <https://doi.org/10.1111/mam.12068>
- Suhonen S, Nummi P, Pöysä H (2011) Long-term stability of boreal lake habitats and use by breeding ducks
- Thompson S, Vehkaoja M, Nummi P (2016) Beaver-created deadwood dynamics in the boreal forest. *Forest Ecology and Management* 360:1–8. <https://doi.org/10.1016/j.foreco.2015.10.019>
- Tickner D, Opperman JJ, Abell R, et al (2020) Bending the Curve of Global Freshwater Biodiversity Loss: An Emergency Recovery Plan. *BioScience* 70:330–342. <https://doi.org/10.1093/biosci/biaa002>
- UNESCO World Water Assessment Programme (2018) The United Nations world water development report 2018: nature-based solutions for water
- Vehkaoja M, Nummi P, Rask M, et al (2015) Spatiotemporal dynamics of boreal landscapes with ecosystem engineers: beavers influence the biogeochemistry of small lakes. *Biogeochemistry* 124:405–415. <https://doi.org/10.1007/s10533-015-0105-4>
- Villéger S, Mason NWH, Mouillot D (2008) New Multidimensional Functional Diversity Indices for a Multifaceted Framework in Functional Ecology. *Ecology* 89:2290–2301. <https://doi.org/10.1890/07-1206.1>

- Vondel BJ van (2005) Haliplidae. In: Nilsson AN, Vondel BJ van (eds) Amphizoidae, Aspidytidae, Haliplidae, Noteridae and Paelobiidae (Coleoptera, Adephaga): (World Catalogue of Insects 7). pp 1–171
- Vondel BJ van (2023) World Catalogue of Haliplidae—corrections and additions. *Koleopterologische Rundschau* 92:73–84
- Washko S, Roper B, Atwood TB (2020) Beavers alter stream macroinvertebrate communities in north-eastern Utah. *Freshwater Biology* 65:579–591. <https://doi.org/10.1111/fwb.13455>
- Washko S, Willby N, Law A (2022) How beavers affect riverine aquatic macroinvertebrates : a review. *PeerJ* 1–19. <https://doi.org/10.7717/peerj.13180>
- Wickham H, Chang W (2016) Package ‘ggplot2’
- Wikar Z, Ciechanowski M, Zwolicki A (2024) The positive response of small terrestrial and semi-aquatic mammals to beaver damming. *Science of The Total Environment* 906:167568. <https://doi.org/10.1016/j.scitotenv.2023.167568>
- Willby NJ, Law A, Levanoni O, et al (2018) Rewilding wetlands: beaver as agents of within-habitat heterogeneity and the responses of contrasting biota. *Philosophical Transactions of the Royal Society B: Biological Sciences* 373:20170444. <https://doi.org/10.1098/rstb.2017.0444>
- Wilson JB, Bremner-Harrison S (2024) A systematic literature review investigating the association between biodiversity and beaver lodges. *Mammal Review* n/a: <https://doi.org/10.1111/mam.12363>
- Wilson KA, Law A, Hester AJ, et al (2024) Tree felling by beaver promotes regeneration in riparian woodlands whilst increasing resource availability for deer. *Forest Ecology and Management* 562:121910. <https://doi.org/10.1016/j.foreco.2024.121910>
- Wright JP, Jones CG, Flecker AS (2002) An ecosystem engineer, the beaver, increases species richness at the landscape scale. *Oecologia* 132:96–101. <https://doi.org/10.1007/s00442-002-0929-1>
- Young TP (2000) Restoration ecology and conservation biology. *Biological Conservation* 92:73–83. [https://doi.org/10.1016/S0006-3207\(99\)00057-9](https://doi.org/10.1016/S0006-3207(99)00057-9)
- Zhang Z, Schwartz S, Wagner L, Miller W (2000) A Greedy Algorithm for Aligning DNA Sequences. *Journal of Computational Biology* 7:203–214. <https://doi.org/10.1089/10665270050081478>

Supporting Information

Table S1. Surveyed wetland details

Wetland number	Wetland type	Wetland name	Latitude	Longitude	Size (Ha)
1	Control	Alaskaivetonjarvet	61.257005	25.119647	1.1
2	Beaver	Downstream Ruuttanajarvi	61.25245	25.09852	2.6
3	Control	Tevajarvet	61.247426	25.09294	0.7
4	Beaver	Saarikonoja	61.22831	25.129314	1.9
5	Beaver	Hauki beaver	61.224267	25.129017	1.7
6	Control	Kylokkaanjarvi	61.2234345	25.1038612	2.5
7	Control	Ylinen Mustajarvi	61.213949	25.10528	1.4
8	Control	Alin Mustajarvi	61.208201	25.114082	0.7
9	Beaver	Peikkoplotti	61.20396	25.135206	0.7
10	Beaver	Peikkoplotti 2	61.2037	25.131316	1.1
11	Beaver	Saukonojan allas	61.200987	25.12862	1.8
12	Beaver	Huhmari 2	61.195086	25.128833	1.2
13	Control	Karvalammi	61.199863	25.1381	0.7
14	Control	Likojarvi	61.199573	25.151788	0.5
15	Control	Syrjanalunen	61.1936223	25.143118	0.9
16	Beaver	Loytjarven	61.1904775	25.1899814	3.8
17	Beaver	Soitimenkorven tekoallas	61.198529	25.19944	4.1
18	Control	Rieskalammi	61.192769	25.206158	2.1

Table S2. A summary of physico-chemical characteristics per wetland type, mean \pm SE (range) with t and p values from linear models that tested differences between each waterbody type.

	Beaver	Control	t and p-value
Size (ha.)	2.1 ± 0.6 (0.7 - 4.1)	1.2 ± 0.2 (0.5 - 2.5)	-2.11, 0.051
Alkalinity (mmol/L)	0.16 ± 0.02 (0.04 - 0.23)	0.08 ± 0.03 (0.00 - 0.23)	-2.14, 0.048*
Chlorophyll A (ug/L)	12.2 ± 4.6 (1.3 - 41.4)	22.8 ± 10.1 (1.1 - 90.1)	0.40, 0.697
Dissolved organic carbon (mg/L)	13.6 ± 2.5 (6.6 - 26.6)	13.6 ± 5.5 (2.3 - 27.2)	-0.23, 0.818
pH	6.3 ± 0.1 (5.4 - 6.8)	5.8 ± 0.2 (4.8 - 6.7)	-1.74, 0.101
Total nitrogen (ug/L)	607.0 ± 88.3 (241.0 - 1021.0)	565.7 ± 79.7 (190.0 - 925.0)	-0.36, 0.722
Total phosphorus (ug/L)	24.8 ± 4.4 (12.0 - 46.0)	27.7 ± 8.5 (7.0 - 93.0)	-0.33, 0.747




Figure S1. Principal components analysis (PCA) of site-level environmental data collected in beaver (blue) and control wetlands (orange). Ellipses cover 95% of the variation in scores for each wetland type on the PCA axes shown.

Table S3. Taxa inventory per taxonomic group per site

[attached excel file]

Table S4. List of indicator species per wetland type for each taxonomic group.

Group	Species	Wetland type	Indicator value	Probability
Aquatic plants	<i>Carex rostrata</i>	Beaver	0.5593	0.001
	<i>Lysimachia thysiflora</i>	Beaver	0.387	0.001
	<i>Hippuris vulgaris</i>	Beaver	0.3287	0.001
	<i>Calamagrostis arundinacea</i>	Beaver	0.2507	0.001
	<i>Lemna minor</i>	Beaver	0.2347	0.001
	<i>Typha latifolia</i>	Beaver	0.2222	0.001
	<i>Potamogeton berchtoldii</i>	Beaver	0.2089	0.001
	<i>Utricularia vulgaris</i>	Beaver	0.1854	0.001
	<i>Cicuta virosa</i>	Beaver	0.1648	0.001
	<i>Utricularia intermedia</i>	Beaver	0.1423	0.001
	<i>Equisetum fluviatile</i>	Beaver	0.1159	0.001
	<i>Utricularia minor</i>	Beaver	0.0995	0.027
	<i>Sparganium glomeratum</i>	Beaver	0.0978	0.001
	<i>Caltha palustris</i>	Beaver	0.0927	0.001
	<i>Carex echinata</i>	Beaver	0.0876	0.001
	<i>Drepanocladus aduncus</i>	Beaver	0.0833	0.003

	<i>Calamagrostis purpurea</i>	Beaver	0.0826	0.001
	<i>Glyceria fluitans</i>	Beaver	0.0774	0.001
	<i>Callitriche hamulata</i>	Beaver	0.0756	0.001
	<i>Scirpus sylvaticus</i>	Beaver	0.0756	0.001
	<i>Scutellaria galericulata</i>	Beaver	0.0692	0.02
	<i>Calliergonella cuspidata</i>	Beaver	0.0633	0.003
	<i>Betula spp.</i>	Beaver	0.0631	0.004
	<i>Agrostis stolonifera</i>	Beaver	0.0554	0.003
	<i>Salix cinerea</i>	Beaver	0.0522	0.05
	<i>Deschampsia cespitosa</i>	Beaver	0.0444	0.004
	<i>Phalaris arundinacea</i>	Beaver	0.0431	0.006
	<i>Salix phylicifolia</i>	Beaver	0.04	0.005
	<i>Sparganium erectum agg.</i>	Beaver	0.04	0.004
	<i>Eleocharis palustris</i>	Beaver	0.0378	0.009
	<i>Alisma plantago-aquatica</i>	Beaver	0.0356	0.013
	<i>Iris pseudacorus</i>	Beaver	0.0356	0.007
	<i>Bidens cernua</i>	Beaver	0.0311	0.023
	<i>Callitriche stagnalis</i>	Beaver	0.0311	0.016
	<i>Juncus filiformis</i>	Beaver	0.0311	0.021
	<i>Equisetum sylvaticum</i>	Beaver	0.0267	0.023
	<i>Lythrum salicaria</i>	Beaver	0.0267	0.033
	<i>Carex lasiocarpa</i>	Control	0.6364	0.001
	<i>Sphagnum auriculatum</i>	Control	0.5385	0.001
	<i>Vaccinium oxycoccos</i>	Control	0.4469	0.001
	<i>Menyanthes trifoliata</i>	Control	0.2995	0.001
	<i>Nuphar lutea</i>	Control	0.242	0.001
	<i>Carex limosa</i>	Control	0.2411	0.001

	<i>Peucedanum palustre</i>	Control	0.2198	0.001
	<i>Phragmites australis</i>	Control	0.2123	0.001
	<i>Sphagnum capillifolium</i>	Control	0.196	0.001
	<i>Andromeda polifolia</i>	Control	0.1805	0.001
	<i>Rhynchospora alba</i>	Control	0.1724	0.001
	<i>Drosera rotundifolia</i>	Control	0.1269	0.001
	<i>Scheuchzeria palustris</i>	Control	0.1156	0.001
	<i>Fontinalis antipyretica</i>	Control	0.1067	0.001
	<i>Rhododendron tomentosum</i>	Control	0.0986	0.001
	<i>Nymphaea candida</i>	Control	0.0918	0.001
	<i>Sphagnum cuspidatum</i>	Control	0.0618	0.011
	<i>Vaccinium uliginosum</i>	Control	0.0515	0.008
	<i>Sphagnum magellanicum</i>	Control	0.0311	0.018
	<i>Carex pauciflora</i>	Control	0.0253	0.047
Beetles	<i>Haliphus ruficollis</i>	Beaver	0.516	0.01
	<i>Hygrotus inaequalis</i>	Beaver	0.499	0.01
	<i>Rhantus exsoletus</i>	Beaver	0.467	0.015
	<i>Hydroporus dorsalis s.s.</i>	Beaver	0.415	0.02
	<i>Psyllobora vigintiduopunctata</i>	Beaver	0.407	0.040
	<i>Ilybius ater</i>	Beaver	0.392	0.025
	<i>Haliphus heydeni</i>	Beaver	0.385	0.015
	<i>Ilybius fuliginosus</i>	Beaver	0.333	0.025
	<i>Hydroporus tristis</i>	Control	0.747	0.005
	<i>Enochrus affinis</i>	Control	0.442	0.02
	<i>Hydroporus neglectus</i>	Control	0.434	0.005
	<i>Hydroporus striola</i>	Control	0.373	0.035
	<i>Contacyphon kongsbergensis</i>	Control	0.363	0.01

	<i>Graphoderus zonatus verrucifer</i>	Control	0.307	0.035
Microcrustacea	<i>Eucyclops</i> spp.	Beaver	0.382	0.015
	<i>Sida</i> spp.	Control	0.384	0.05
True flies	<i>Simulium</i> spp.	Beaver	0.597	0.005
	<i>Conchapelopia</i> spp.	Beaver	0.516	0.005
	<i>Phaenopsectra</i> spp.	Beaver	0.509	0.005
	<i>Microtendipes</i> spp.	Beaver	0.479	0.005
	<i>Psectrotanypus</i> spp.	Beaver	0.461	0.015
	<i>Stempellinella</i> spp.	Beaver	0.416	0.005
	<i>Anopheles</i> spp.	Beaver	0.403	0.01
	<i>Eukiefferiella</i> spp.	Beaver	0.376	0.005
	<i>Monopelopia</i> spp.	Beaver	0.376	0.005
	<i>Nanocladius</i> spp.	Beaver	0.376	0.005
	<i>Diamesa</i> spp.	Beaver	0.375	0.01
	<i>Arctopelopia</i> spp.	Beaver	0.36	0.005
	<i>Dicrotendipes</i> spp.	Beaver	0.36	0.005
	<i>Rheocricotopus</i> spp.	Beaver	0.341	0.01
	<i>Tanypus</i> spp.	Beaver	0.305	0.01
	<i>Prionocera</i> spp.	Beaver	0.303	0.005
	<i>Notiphila</i> spp.	Beaver	0.244	0.04
	<i>Tipula</i> spp.	Beaver	0.24	0.015
	<i>Trissopelopia</i> spp.	Beaver	0.24	0.04
	<i>Bryophaenocladus</i> spp.	Control	0.335	0.010
	<i>Smittia</i> spp.	Control	0.282	0.025
May/Stone/Caddisflies	<i>Leuctra</i> spp.	Beaver	0.547	0.025
Fishes	<i>Carassius carassius</i>	Control	0.1176	0.005




Figure S2. Error plots comparing taxa richness per sample for beaver (blue) and control wetlands (orange) for a) aquatic plants, b) beetles, c) segmented worms, d) microcrustacea, e) true flies, f) may/stone/caddisflies, g) amphibians, h) fish, i) birds and j) mammals. Coloured points represent the mean of the data, error bars are equal to one standard deviation, and dots show individual data points. Z test statistics and p values are outputs from generalised linear mixed models that tested differences between waterbody types per taxonomic group.




Figure S3. Error plots comparing taxa richness per site for beaver (blue) and control wetlands (orange) for a) aquatic plants, b) beetles, c) segmented worms, d)

Figure S4. Error plots comparing site occupancy (presence across 10 samples) of aquatic plant species between beaver (blue) and control (orange) wetlands. Coloured points represent the mean of the data, error bars are equal to one standard error and dots show individual data points. P values are outputs from Benjamini-Hochberg corrected non-parametric Wilcoxon tests that tested differences between waterbody types per genera.




Figure S5. Error plots comparing site occupancy (presence across 10 samples) of beetle species between beaver (blue) and control (orange) wetlands. Coloured points represent

the mean of the data, error bars are equal to one standard error and dots show individual data points. P values are outputs from Benjamini-Hochberg corrected non-parametric Wilcoxon tests that tested differences between waterbody types per genera.




Figure S6. Error plots comparing site occupancy (presence across 10 samples) of segmented worm genera between beaver (blue) and control (orange) wetlands. Coloured points represent the mean of the data, error bars are equal to one standard error and dots show individual data points. P values are outputs from Benjamini-Hochberg corrected non-parametric Wilcoxon tests that tested differences between waterbody types per genera.




Figure S7. Error plots comparing site occupancy (presence across 10 samples) of microcrustacea genera between beaver (blue) and control (orange) wetlands. Coloured points represent the mean of the data, error bars are equal to one standard error and dots show individual data points. P values are outputs from Benjamini-Hochberg corrected non-parametric Wilcoxon tests that tested differences between waterbody types per genera.




Figure S9. Error plots comparing site occupancy (presence across 10 samples) of may/stone/caddisfly genera between beaver (blue) and control (orange) wetlands. Coloured points represent the mean of the data, error bars are equal to one standard error and dots show individual data points. P values are outputs from Benjamini-Hochberg corrected non-parametric Wilcoxon tests that tested differences between waterbody types per genera.




Figure S10. Error plots comparing site occupancy (presence across 10 samples) of amphibian species between beaver (blue) and control (orange) wetlands. Coloured points represent the mean of the data, error bars are equal to one standard error and dots show individual data points. P values are outputs from Benjamini-Hochberg corrected non-parametric Wilcoxon tests that tested differences between waterbody types per species.




Figure S11. Error plots comparing site occupancy (presence across 10 samples) of fish species between beaver (blue) and control (orange) wetlands. Coloured points represent the mean of the data, error bars are equal to one standard error and dots show individual data points. P values are outputs from Benjamini-Hochberg corrected non-parametric Wilcoxon tests that tested differences between waterbody types per species.




Figure S12. Error plots comparing site occupancy (presence across 10 samples) of bird species between beaver (blue) and control (orange) wetlands. Coloured points represent the mean of the data, error bars are equal to one standard error and dots show individual data points. P values are outputs from Benjamini-Hochberg corrected non-parametric Wilcoxon tests that tested differences between waterbody types per species.




Figure S13. Error plots comparing site occupancy (presence across 10 samples) of mammal species between beaver (blue) and control (orange) wetlands. Coloured points represent the mean of the data, error bars are equal to one standard error and dots show individual data points. P values are outputs from Benjamini-Hochberg corrected non-parametric Wilcoxon tests that tested differences between waterbody types per species.

eDNA Methods

Field Sampling and eDNA Capture

Shoreline water samples were collected for eDNA analysis from all 18 wetlands. Each individual sample consisted of 1.5 L of surface water collected from five points within a roughly 20 m zone, parallel to the shoreline. Samples were collected at 10 roughly equidistant points, determined by accessibility, around the perimeter of each wetland in sterile Whirl-Pak® bags, resulting in 180 samples in total. One 1.5 L bag of UV filtered and distilled water was carried into the field alongside samples for each wetland. Samples and blanks were placed in ice-filled insulated cool boxes lined with site-specific sterile plastic bag liners. Fresh gloves were used for each sample. Samples were transported to the field station for filtering. Lab surfaces were sterilised using 10% bleach solution before filtering. During filtering, sterile equipment was kept out of contact with contaminated surfaces and equipment and gloves replaced between samples. All samples and field blanks, were manually pumped through 0.45 µm PVDF Sterivex filter units (micrometer), using sterile 60 ml syringes until the filter clogged, then the filtered volume was recorded. Volumes ranged from 70 to 750 ml between samples. Air was then forced through the filter to remove any excess liquid water from the capsule. Each Sterivex filter was then filled with 1 ml of pre-aliquoted Longmires buffer solution (Longmire et al. 1997) before sealing with a screw cap and parafilm. Filter units were stored at room temperature at Lammi and -20°C on arrival in the UK.

DNA Extraction, Library Preparation and Sequencing

DNA was extracted from Sterivex filters using a modified version of the Modular Universal DNA (MuDNA) extraction protocol (Sellers et al. 2018) in a dedicated eDNA lab at the University of Hull. The Longmire's solution was pushed from the filter into a 2 ml eppendorf tube and centrifuged at 6000 g for 30 minutes. Supernatant was discarded and the pellet resuspended in 60 μ l of Lysis Solution (Sellers et al. 2018) by vortexing. To remove the remaining DNA from the filter, a mixture of 660 μ l of Lysis Solution, 180 μ l 6% SDS and 60 μ l 10 mg/ml Proteinase K was added, and placed in a rotating incubator overnight at 55°C. The Inhibitor Removal Step of the MuDNA: Water protocol was then followed. The lysis mixture was pulled from the filter unit and 300 μ l of Flocculant Solution (0.3x volume, Sellers *et al.* (2018)) added before vortexing and incubating on ice for 10 minutes, centrifuging at 10,000 g for 2 minutes and transferring the supernatant to a new 2 ml tube. The Silica Binding and Wash steps of the MuDNA protocol: Water were followed as in Sellers *et al.* (2018), but the Wash step was repeated once more. DNA was eluted in 200 μ l Elution Buffer (Sellers et al. 2018). Extraction blanks of sterile Longmire's buffer solution in an unused Sterivex filter were processed under the same conditions to detect any contamination during the extraction stage. eDNA samples were stored at -20°C.

DNA extractions were PCR-amplified in two-step protocols. Vertebrate-specific primers that amplify a 106 bp fragment of the mitochondrial 12S ribosomal RNA region (Riaz et al. 2011; Kelly et al. 2014), and primers that target a 141bp region of the COI in benthic invertebrates (fwhF2/EPTDr2n) (Leese et al. 2021) were used.

For vertebrates, eight nucleotide long Multiplex Identification (MID) Tags were included in both first and second round PCRs for unique sample identification following a nested tagging approach described in (Kitson et al. 2019). Twenty-four unique MID tags were used for the forward and another 24 for the reverse primers. First round PCR primers

also included heterogeneity spacers, sequencing primers and pre-adapters. Samples were processed in sub-libraries consisting of 24 samples including one field (filtration) blank, one extraction blank, one PCR negative and one positive control (DNA from the African cichlid, zebra mbuna, *Maylandia zebra*, at 0.05 ng/ μ l). First round PCRs were performed in triplicate, in final reaction volumes of 25 μ l, with 12.5 μ l of Q5[®] High-Fidelity 2x Master Mix (New England Biolabs), 0.5 μ l of Thermo Scientific Bovine Serum Albumin (Fisher Scientific UK Ltd.), 7 μ l of Molecular Grade Water (Fisher Scientific UK Ltd.), 1.5 μ l of each 10 μ M tagged primer, and 2 μ l of template DNA. PCR were performed in a Veriti Thermocycler (Applied Biosystems) with the following profile: 98°C for 5 mins, 35 cycles of 98°C for 10 s, 58°C for 20 s and 72°C for 30 s, 72°C for 7 mins then held at 4°C. Triplicate PCR products were pooled, and 2 μ l run on 2% agarose gels to check amplification. Sub-libraries were then pooled according to band strength (no/very faint band = 20 μ l, faint band = 15 μ l, bright band = 10 μ l, very bright band = 5 μ l) on a gel (Alberdi et al. 2018) before a double-size selection bead purification step. Ratios of 0.9x and 0.15x Mag-BIND RxnPure Plus magnetic beads (Omega Bio-tek) to 100 μ l of each sub-library were used for purification, and sub-libraries eluted in 25 μ l Elution Buffer (1xTE).

Two replicates of each sub-library were performed for second round PCR, with one unique forward and one unique reverse MID tag per sub-library (i.e. 10 in total). Second round PCRs were carried in 50 μ l reaction volumes, with 25 μ l of Q5[®] High-Fidelity 2x Master Mix (New England Biolabs), 13 μ l of Molecular Grade Water (Fisher Scientific UK Ltd.), 3 μ l of each 10 μ M tagged primer, and 4 μ l of template DNA. PCRs were carried out on a Veriti Thermocycler (Applied Biosystems) with the following thermocycling profile: 95°C for 3 mins, 10 cycles of 98°C for 20 s and 72°C for 1 min, 72°C for 5 mins then held at 4°C. Duplicates were pooled and PCR products checked on a 2% agarose gel before a second

bead clean, with ratios of 0.7x and 0.15x Mag-BIND RxnPure Plus magnetic beads (Omega Bio-tek) to 50 μ l of each sub-library. Sub-libraries were eluted in 25 μ l and quantified on a Qubit™ 3.0 fluorometer using a dsDNA HS Assay Kit (Invitrogen) then normalised by pooling according to sample size and library concentration, and bead cleaned as for the PCR2 protocol. The library was diluted to 4 nM based on Qubit™ concentration, then quantified by qPCR using the NEBNext Library Quant Kit for Illumina (New England Biolabs) on a StepOne Thermocycler (Applied Biosystems). The final library was adjusted to 4 nM based on qPCR results and denatured and diluted according to the Illumina MiSeq library preparation guide. The final libraries were sequenced at 13 pM on an Illumina MiSeq using 2 x 300 bp V3 chemistry (Illumina).

To amplify invertebrate data for sequencing, the following changes were made to library preparation methods. No MID tags were included in PCR 1 where QIAGEN Multiplex PCR Plus Kits were used in place of the Master Mix. DNA extracted from *Calliphora vicina* was used as the positive control. Respective PCR conditions as follows; 10 cycles of 95°C for 30 s, 64°C for 90 s (-1°C each cycle), 72°C for 30 s. 25 cycles of 95°C for 30 s, 54°C for 90 s, 72°C for 30 s, a final elongation of 68°C for 10 mins and finally held at 4°C. Pooled PCR 1 triplicates were then purified with 0.9x vortex suspended magnetic beads per sample. PCR 2 commenced immediately after confirming the PCR 1 reactions and bead cleans were successful using a 2% agarose gel. MID tags were introduced at this stage, with 240 sample specific combinations of 24 reverse and 10 forward sequences. Product from PCR 1 was tagged and further amplified in 25ul volumes per duplicate reaction well for each sample. Using Q5 High-Fidelity 2x Master Mix (New England Biolabs) and 0.5 μ l more water instead of Bovine Serum. Invertebrate PCR 2 conditions are as per vertebrate PCR 2 conditions. After pooling duplicates and samples into sub-libraries by agarose gel band strength, a final

magnetic bead purification is completed with a ratio of 0.9x suspended mixture to sample. All subsequent library preparation steps are identical to the previous vertebrate sequencing methods.

Bioinformatics

For vertebrates, a custom reference database of published 12S rRNA sequences, downloaded from GenBank, was created based on records of the Finnish Biodiversity Information Facility, FinBIF (<https://laji.fi/en>) and local expert knowledge of staff from the University of Helsinki Lammi Field Station on the vertebrate biodiversity of the Evo region. Poor quality sequences (those with > 100 ambiguous bases) were removed from the reference database. For invertebrates an updated version of the reference database published by Harper et al. (2019a) was used.

Vertebrate sequencing data was demultiplexed to forward and reverse fastq files per library using Illumina MiSeq Reporter software, then to sample using a custom Python script. Further quality control and taxonomic assignment for both libraries were performed using Tapirs (<https://github.com/EvoHull/Tapirs>), a Snakemake workflow manager (Köster and Rahmann 2012) for reproducible analysis of DNA metabarcoding data. Quality control included trimming, denoising and dereplication. Raw reads were quality trimmed from the tail with a 5 bp sliding window (qualifying phred score of Q30 and an average window phred score of Q30) using fastp (Chen et al. 2018), allowing no more than 40% of the final trimmed read bases to be below Q30. Primers were removed by trimming the first 18 bp of forward and reverse reads. For vertebrates and invertebrates respectively reads were tail cropped to a maximum length of 106 bp and 142 bp and reads shorter than 85bp and 100 bp were

discarded. Sequence read pairs were merged into single reads using fastp, provided there was a minimum overlap of 20 bp, no more than 5% mismatches and no more than 5 mismatched bases between pairs. Only forward reads were kept from read pairs that failed to be merged. A final length filter removed any reads longer than 110 bp for vertebrates and 160 bp for invertebrates to ensure sequence lengths approximated the expected fragment size (~106 bp or ~ 142 bp respectively).

Redundant sequences were removed by clustering at 100% read identity and length (--derep_fulllength) in VSEARCH (Rognes et al. 2016). Clusters represented by less than three sequences were omitted from further processing. Reads were further clustered (--cluster_unoise) to remove redundancies due to sequencing errors (retaining all cluster sizes). Retained sequences were screened for chimeric sequences with VSEARCH (--uchime3_denovo).

The final clustered, non-redundant query sequences were then compared against the reference database using BLAST (Zhang et al. 2000). Taxonomic identity was assigned using a custom majority lowest common ancestor (MLCA) approach based on the top 2% query BLAST hit bit-scores, with at least 90% query coverage and a minimum BLAST hit identity of 98% for vertebrates and 95% for invertebrates. Of these filtered hits, 80% of unique taxonomic lineages therein had to agree at descending taxonomic rank (domain, phylum, class, order, family, genus, species) for it to be assigned a taxonomic identity. If a query had a single BLAST hit it was assigned directly to this taxon only if it met all previous MLCA criteria. Read counts assigned to each taxonomic identity were calculated from query cluster sizes. Lowest taxonomic rank was to species and assignments higher than order were classed as unassigned. For vertebrates, reads from human, positive control (cichlid),

unassigned, domestic species (*Canis* spp., *Felis* spp., *Sus scrofa*, *Bos taurus*, *Meleagris gallopavo*), and species that have not been previously recorded in the study area (*Alburnus alburnus*, *Cobitis taenia*, *Cottus gobio*, *Cyprinus carpio*, *Gasterosteus aculeatus*, *Gobio gobio*, *Lota lota*, *Microtus rossiaemeridionalis*, *Pungitius pungitius*, *Rhodeus amarus*, *Salmo salar*, *Salmo trutta*) were removed from the dataset for downstream analyses. The genera *Aythya* and *Mareca* were assigned to Anatidae and *Columba livia* to Columbidae. For invertebrates, unassigned and the positive *Calliphora vicina* were removed.

eDNA raw data and quality control

For vertebrates, after demultiplexing to sample, raw data consisted of 21,540,228 paired end reads across all samples and controls. Following taxonomic assignment against our curated reference database, 16,729,221 reads remained. 53 OTUs were retained for the final dataset. After quality control steps including application of the contamination threshold and removal of non-target species, 175 samples remained, averaging 10,444 reads per sample.

For invertebrates, 14,818,440 reads remained after taxonomic assignment. A Limits Of Detection (LOD) approach was used to apply a sample-specific contaminate threshold of the mean plus standard deviation*3 of all reads from taxa detected in the Negative and Positive PCR controls (mean + SD*3). This framework was then adjusted for field and extraction controls to use the max (instead of mean) reads of any taxa found across these blanks (max+SD*3). 177 samples remained, averaging 3216 reads per sample after thresholds were applied.